

The influence of androgenic steroid hormones on female aggression in 'atypical' mammals

Jeffrey A. French, Aaryn C. Mustoe, Jon Cavanaugh and Andrew K. Birnie

Phil. Trans. R. Soc. B 2013 **368**, 20130084, published 28 October 2013

References

[This article cites 90 articles, 16 of which can be accessed free](#)

<http://rstb.royalsocietypublishing.org/content/368/1631/20130084.full.html#ref-list-1>

Subject collections

Articles on similar topics can be found in the following collections

[behaviour](#) (462 articles)

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Review



Cite this article: French JA, Mustoe AC, Cavanaugh J, Birnie AK. 2013 The influence of androgenic steroid hormones on female aggression in 'atypical' mammals. *Phil Trans R Soc B* 368: 20130084. <http://dx.doi.org/10.1098/rstb.2013.0084>

One contribution of 14 to a Theme Issue 'Female competition and aggression'.

Subject Areas:

behaviour

Keywords:

aggression, dominance, testosterone, organizational effects, activational effects, mammals

Author for correspondence:

Jeffrey A. French
e-mail: jfrench@unomaha.edu

The influence of androgenic steroid hormones on female aggression in 'atypical' mammals

Jeffrey A. French^{1,2}, Aaryn C. Mustoe¹, Jon Cavanaugh¹ and Andrew K. Birnie¹

¹Department of Psychology, Callitrichid Research Center, and ²Department of Biology, University of Nebraska at Omaha, Omaha, NE 68182, USA

Dimorphism on dominance and agonistic behaviour in mammals tends to be strongly biased toward males. In this review, we focus on a select few species of mammals in which females are as or more aggressive than males, and/or are dominant to males, and explore the role of androgenic hormones in mediating this important difference. While the data are not as clear-cut as those published on traditional laboratory mammals, our review highlights important endocrine substrates for both organizational and activational influences of steroids on female aggressive behaviour. We highlight areas in which further observations and experiments are crucial, especially the potential facilitative effects of androgens on female aggression. Finally, new and innovative techniques, including molecular genetics and receptor pharmacology, portend important insights into the ways in which androgenic hormones regulate aggressive behaviour in 'atypical' female mammals.

1. Introduction

When one views the approximately 5000 species in Class Mammalia writ large, few would contest the statement that, among the species in this class, males are more aggressive than females. This truism is reflected in prominent class-wide sexual dimorphisms in the rate of aggressive interactions, in differences in time and energy budgets allocated to aggressive and dominance interactions, variation between the sexes in the proportion of aggressive elements in the behavioural repertoire and in the evolution of traits (e.g. large body size or morphological adaptations for fighting such as large canines, horns and antlers [1,2]). Each of these traits facilitates aggressive competition among males. Traditional explanations for this prominent pattern of mammalian sexual dimorphism, vis-à-vis aggressive behaviour and its associated morphological weaponry, rely primarily on the impact of sexual selection for male traits that facilitate or promote intrasexual competitive ability, and hence reproductive success [3]. However, recent analyses have strongly suggested that this explanation is neither exhaustive nor definitive (cf. [4,5]).

As with all biological 'truisms', however, there are notable exceptions to the general rule, and exploring the causes and consequences of female aggression in species where females are as aggressive or even more aggressive than males can provide important insights into the proximate and ultimate mechanisms that underlie these differences. Ralls [6] was among the first to bring to the attention of biologists those curious cases of species in which typical mammalian sexual dimorphism in size did not hold true. This work has stimulated detailed analyses of the phylogeny, evolution and ecological contexts that are associated with mammalian species in which females are, on average, the same size or larger than males, and research on other species in which females are as or more aggressive than males (see reviews in [7,8]; [9]). To the extent that selection has favoured female aggression among mammals, one would expect to see 'fingerprints' of this selection in the underlying proximate physiological mechanisms that promote and facilitate aggression in females. Given the conservative nature of evolution, we would expect that selection for female aggression

would operate on common or similar neurobiological pathways that are well known to regulate both enhanced male aggression (relative to females) and individual differences among males in the display of aggressive behaviour.

Given the importance of androgenic steroid hormones in both the ontogenetic organization of neural structures underlying sex-typical behaviour in males, including aggression [10], and in activating or facilitating aggressive behaviour in adult males [11,12], we focus on the impact of androgens on female aggressive phenotypes. Our goal in this paper is to highlight historical and current research on the proximate neuroendocrine mechanisms underlying female aggression in a select group of mammals. Criteria for selection of these species are twofold: first, we selected species in which mammalian-typical male-biased aggression is either absent or even sex-reversed. We included cases in which intrasexual aggression in females is conspicuous and also cases where females are more aggressive than males, and generally win fights with male conspecifics and/or exhibit dominance over males. Second, we limited our review to species for which published data on the impact of gonadal steroids on female aggression are sufficiently detailed to allow a careful assessment of the mechanisms that contribute to this atypical (for mammals) pattern.

2. Spotted hyaena (*Crocuta crocuta*)

Ever since Aristotle's formal writings over 2000 years ago on the observation that 'every [spotted] hyaena is furnished with the organ both of the male and the female' (cited in [13, p. 349]), the attention of biologists has been drawn to this species as a model for endocrine substrates for sexual differentiation of genital morphology. The conspicuously high levels of aggression in adult females and almost universal female dominance over males in clan societies also provide an exemplary natural model for exploring the differentiation of aggressive behaviour and its endocrine correlates. There is a vast and contentious literature on the ontogenetic origins of genital differentiation (which will certainly make an interesting study for historians of science; e.g. [13]), but for the purposes of this review, we limit our discussion to the role of endocrine exposure at distinct developmental stages and the link to aggressive behaviour and dominance. Below we briefly explore the evidence for differential prenatal, early postnatal and adult hormone milieus in hyaenas, and their implications for sex differences in the brain nuclei associated with social behaviour, and aggression and dominance within hyaena clans.

(a) Prenatal hormone exposure

In spotted hyaenas, levels of sex steroids rise throughout pregnancy in adult females, with levels of androgens (testosterone: T; androstenedione: A₄) peaking in the last trimester of gestation [14]. A high placental level of the steroidogenic enzyme 17 β -HSD converts A₄ to T, providing a rich uterine environment with potent bioactive androgens [14,15]. This gestational androgen environment would appear to set the stage for potential masculinization of female cubs.

The most detailed evidence that prenatal hormone exposure alters patterns of postnatal aggressive behaviour comes from analyses of individual differences in maternal androgens and later-life behaviour in wild hyaenas. Dloniak *et al.* [16] confirmed the general observation of elevated maternal androgens

in pregnancy, particularly in the second half of gestation. Androgen levels in this period, during which behavioural masculinization is most likely, were higher in dominant females than in subordinates. Female and male cubs exposed to higher levels of maternal androgen in the second half of pregnancy had higher levels of aggression and male-like sexual mounting than cubs exposed to lower levels of androgens, even when controlling for maternal rank and gestational glucocorticoid exposure [17]. Further analyses of lifetime rates of aggression in females revealed that rates of intersexual aggression were significantly and positively associated with gestational androgen exposure, but intrasexual aggression by females was not related to prenatal androgen exposure [18]. These data strongly implicate late-pregnancy androgen exposure as a critical determinant of both juvenile and adult female aggression.

A series of studies on the spotted hyaena have addressed organizational effects of androgens by administering a potent androgen receptor blocker (flutamide) and an aromatase inhibitor (finasteride) to pregnant females, a treatment regimen that effectively minimizes exposure of fetal hyaenas to androgenic hormones. When sampled as adults, both male and female hyaenas in the prenatal antiandrogen treatment had lower levels of T, higher levels of oestradiol (E₂) and an enhanced (i.e. more 'feminine') luteinizing hormone response to gonadotropin-releasing hormone stimulation [19]. There is an unpublished observation that prenatal antiandrogen treatment alters aggressive behaviour in females, in the form of reduced female–female neonatal sibling fights (cited in [20]), fights that are common and quite often fatal [21,22].

(b) Prepubertal hormone exposure

Female and male spotted hyaenas experience highly variable steroid hormone environments as they approach puberty. Glickman *et al.* [23] documented normative levels of sex steroids in male and female hyaenas from two to 60 months of age, and interesting sex differences emerged. First, and not surprisingly, levels of E₂ rose throughout prepubertal development in females, but not in males. Early in postnatal life, T levels did not differ between males and females, but levels increased in males beginning at 20 months of age. By 60 months of age, male T levels were five to sixfold higher than those in females. Conversely, A₄ levels showed the reverse pattern: levels were higher in females than in males beginning early in life and these sex differences persisted up to 60 months of age. Although these normative patterns of steroid exposure have not been explicitly linked with either subadult or adult aggression, removing the source of the hormones in females alters both steroid and behavioural characteristics. Neonatally ovariectomized (OVX) females displayed low or undetectable levels of both T and A₄, confirming the ovary as the source of these male-typical sex steroids in females [23]. An unpublished study implicates variation in postnatal steroid levels and aggressive behaviour in females: prepubertal ovariectomy reduced adult female aggression toward males (cited in [13]). These results are consistent with a postnatal endocrine influence on female aggression, but stronger tests of this hypothesis are required.

(c) Hormones and aggression in adult females

There is consensus in the literature that T levels in adult males (regardless of whether they live in their natal clan or

have dispersed to a new clan) are higher than those in adult females; thus, female sex-biased aggression and dominance cannot be directly traced to elevated T [14,16,24]. By contrast, levels of the important T precursor, A_4 , are similar in adult male and female hyaena [14,24]. Without the knowledge of activity of the enzyme 17 β -HSD that converts A_4 to T, and more importantly, sex differences in the bioactivity of this enzyme, it is impossible to determine whether T derived from A_4 conversion could potentially account for the prominent atypical sex difference in aggression and dominance in spotted hyaenas. Studies of brain structure in adult spotted hyaenas reveal departures from more traditional mammals that hint at neural substrates for enhanced aggression. The medial preoptic area and anterior hypothalamus play important roles in mediating male-typical behaviour in mammals, and the sexually dimorphic nucleus of this region, normally substantially larger in male than female mammals, was only slightly smaller in female hyaenas than in males [25]. Further, in typical mammals, males have higher densities of vasopressin-containing neurons in the lateral septum (which mediates aggressive behaviour), but in a small sample of adults, female hyaenas tended to have more vasopressin neurons than males [26]. In spite of the potential for steroids (particularly A_4) to serve an activating role for female aggression, and neural structures that are consistent with a female brain primed for aggressive behaviour, there has yet to be a good empirical test of the activational role of androgens in female spotted hyaenas. Although a facilitative effect of engaging in high levels of agonistic behaviour in elevating future levels of androgens has not been explicitly addressed as of yet in females, levels of androgens in male hyaenas are more strongly predicted by their rates of interaction with more aggressive and dominant females than by rates of male–male aggression [27]. Exploring the extent to which androgen levels in adult females vary as a consequence of rates of agonistic interactions, and subsequently alter thresholds for future aggression, would thus appear to be a promising area for additional research.

More recent evidence on spotted hyaena biology highlights the notion that circulating levels of hormones *alone* may not explain the masculinization of female genital morphology and the sex-reversed pattern of aggressive behaviour and dominance in this species. Browne *et al.* [28] report that the histomorphology of male and female gonads are distinct as early as day 30 of gestation (GD30), with male gonads expressing anti-Mullerian hormone (AMH) and three enzymes crucial for androgen biosynthesis (P450c17, cytochrome b5 and 3 β -HSD). By contrast, female gonads at GD30 lacked AMH and all three enzymes. However, by GD95, ovarian tissue displayed immunohistochemical staining for all three androgen-synthesizing enzymes. Thus, there appear to be two different windows for fetal androgen synthesis for males and females. In males, the enzymatic substrates are available for androgen synthesis as early as GD30, potentially contributing to masculinization of male genitalia early in development and brain/behaviour later in gestational development. For females, by contrast, the lack of substrates for androgen synthesis in the ovary at GD30 may facilitate the growth and elaboration of accessory reproductive structures along a ‘female’ path (e.g. development of urethral retractor muscles and the pleating and elasticity of the ‘future’ birth canal). The presence of the ovarian androgen-synthesizing enzymes later in gestational development (GD90) could provide the endocrine substrates

for neural masculinization (see above) underlying both neonatal and adult aggression in female hyaenas.

There are also important differences in carrier proteins for steroid hormones in the spotted hyaena that may contribute to the unique process of sexual differentiation in this species (box 1). This carrier protein (sex hormone-binding globulin; SHBG) is critical not only for increasing the half-life of steroid hormones in circulation, but also renders steroids biologically inactive while they are bound to it. Gene sequences for SHBG in spotted hyaenas were contrasted with those in three closely related species that display mammal-typical patterns of male dominance and aggression and dimorphic external genitalia [36]. Relative to the ‘typical’ hyaenids, spotted hyaenas have two missing amino acids and a single amino acid substitution in the SHBG secretion signal polypeptide. Modification of the spotted hyaena SHBG gene to resemble the structure of other hyaenid genes increases SHBG production, suggesting that the molecular differences in the spotted hyaena SHBG gene reduce SHBG production (and hence increase the availability of bioactive steroids to act on target tissues). Further, Hammond *et al.* [36] demonstrated that while females in the typical hyaenids have substantially higher titres of SHBG than males (and hence are ‘protected’ from excess androgen exposure), female spotted hyaenas had low circulating levels of SHBG, which were not statistically distinguishable from those of males. Thus, mutations in the genes regulating SHBG in spotted hyaenas, relative to other hyaenids, appear to enhance exposure of females to elevated levels of bioactive steroids, including androgens of ovarian origin, and these steroid carrier proteins may, therefore, be important contributors to variation in steroid hormone exposure.

3. Lemurs (*Lemur spp.*)

The ring-tailed lemurs (*Lemur catta*) serve as a fascinating model to explore proximate mechanisms for female aggression. Female lemurs are unique in that they exhibit unconditional dominance over adult males and sometimes over other females. This female-dominant social structure in ring-tailed lemurs was first reported by Jolly [37], and patterns of female dominance were later described in other lemur species [38–40]. Adult male lemurs will exhibit spontaneous submissive behaviours toward mature female conspecifics [41,42] and this relationship of female dominance over males occurs in reproductively mature lemurs only [43]. Moreover, females are generally intolerant of other conspecifics and fail to provide support to others during conflicts [44]. Nearly all lemurs are seasonal breeders and aggression peaks during times of peak mating and birth seasons [45], and female lemur aggression is intimately linked with patterns of feeding [46,47].

Male-like traits in female lemurs are evident in their genital morphology. Female lemurs possess masculinized genitalia including an elongated, enlarged clitoris [48], and there are potential indications that this ‘masculinization’ could be influenced by fetal androgen exposure [49]. For instance, lemurs pregnant with female offspring possess higher androgen/oestrogen ratios, and, consequently, the relative and not absolute values of prenatal androgens may be partially responsible for the masculinization seen among female lemurs [50]. While the endocrinology of pregnancy in female ring-tailed lemurs seems appropriate for the

Box 1. Issues in the measurement of androgens in female mammals.

There are several methodological issues that must be considered when measuring steroid hormones in female mammals, some of which are unique to females and others that are equally applicable to males and females.

Source of hormone. Multiple tissues in females are capable of synthesizing and secreting androgen hormones, including the ovary, placenta, fetus and adrenal cortex. While ovaries are the primary sources of oestrogens and progestins, thecal cells have the potential to produce androgens and, depending upon ovarian levels of aromatase, are either converted to oestrogens or secreted as androgens [29]. Conceptive ovarian cycles in a number of mammalian species, including dogs, marmosets, baboons and humans, are characterized by elevated androgens likely of luteal origin (see references in Fite *et al.* [30]). The placenta is a steroid-rich tissue, and androgens and their metabolic precursors can be produced from this source [14]. Testicular production of androgens by male fetuses can cross the fetal–placental barrier and enter general circulation in pregnant females, although this does not inevitably occur in all species (literature reviewed in French *et al.* [31]). Finally, androgens of adrenal origin, although typically secreted at substantially lower levels than those of gonadal origin, can nonetheless contribute to overall circulating androgen levels [32].

Biological substrates. Traditionally, behavioural endocrinologists have relied on measurements of circulating androgens in plasma or serum, as circulating hormones are those that have the opportunity to interact with target tissues, such as steroid-sensitive neuronal circuits in the brain. Non-invasive sampling regimens have been developed in both laboratory and especially field contexts, and steroids are now measured in a variety of biological matrices, such as urine, faeces, hair, saliva and milk. This creates two issues for the interpretation of concentrations. First, clearance rates for steroids differ depending upon the biological sample, in the order of minutes in saliva, minutes-to-hours in urine and hours-to-days in faeces [33]. Quantifying the clearance rates of labelled steroids can clarify these temporal issues. Second, significant metabolism of steroids occurs from secretion to excretion, particularly in the liver [34], thus the hormone measured in excreta can differ from the hormone acting on target tissues. Again, administration of labelled hormone and characterization of the chemical structure of the labelled excreted hormone can identify important metabolic transformation of the hormone of interest.

Cross-reactivity of antibody. As with any immune-based quantitative assay, immunoassays for androgenic steroids use antibodies of varying specificity for steroids. Antibodies with high specificity (low cross-reactivity) for a specific steroid provide the confidence that the resulting quantitative measures are selective for that steroid, but antibodies with higher cross-reactivity with chemically similar steroids (and hence less selectivity) provide quantitative data that represent the sum of multiple steroid concentrations, adjusted by the degree of cross-reactivity for steroids other than the target hormone against which the antibody has purportedly been generated. For investigators interested in the potential actions of specific steroids on behavioural outputs, antibodies with high specificity or sample purification with chromatographic separation are essential. On the other hand, antibodies with high cross-reactivity can provide information not only on the hormone of interest, but also on important levels of precursors and biologically active downstream metabolites of the hormone of interest. For example, Dloniak *et al.* [16] used an antibody directed against testosterone to measure androgens in spotted hyaena faecal samples. Chromatographic separation of androgens by polarity and subsequent immunoassay with the T antibody revealed high cross-reactivity with both androstenedione (an important prohormone for T) and dihydrotestosterone (a biologically active metabolite of T; [16]). Thus, the lack of high specificity of the T antibody in fact provides a more integrative measure of potential and available steroid levels. Clarity and caution regarding the resulting measures in these cases is required: the concentrations are accurately expressed as ‘faecal androgens’ as opposed to ‘faecal T’.

Carrier proteins. Steroids often circulate in the blood while bound to glycoproteins known as sex hormone-binding globulin (SHBG). In the event that steroids are complexed with SHBG, they are incapable of crossing cell membranes, do not interact with nuclear steroid hormone receptors, and hence do not trigger transcriptional changes in the nucleus, which is the primary mechanism of steroid hormone action [35]. Quantification of the per cent of circulating androgens that are bound to SHBG or are circulating in a free form is crucial in interpreting the potential pool of bioactive hormones that can interact with and alter brain function.

behavioural masculinization of female fetuses (e.g. high levels in the third trimester of pregnancy, regardless of litter sex ratio [49]), specific organizational effects of androgens on behavioural profiles have yet to be demonstrated.

Given the strong and pervasive link of aggression and dominance in male mammalian species and increased androgen production [12,51], it seems likely that female lemurs with increased T levels might engage more frequently in dominant and aggressive behaviours. While a wealth of data are available for evaluating the organizational and activational hormonal contexts of female dominance found among spotted hyaenas (*Crocuta*, discussed in this article), only a few studies have examined the relationship between female dominance,

aggression and androgen production in female lemurs [52,53]. Both aggression and circulating T in females peak during the lemur breeding season [45,54] and, consequently, it is easy to speculate that circulating T (and other androgens) may be facilitating the breeding season peak in female aggressive behaviour. Von Engelhard *et al.* [53] found that while both faecal T and female aggression in lemurs increased twofold during the breeding season, individual female aggression rates and individual indices of female dominance were not correlated with androgen concentrations. Additionally, female lemurs did not have higher levels of androgens relative to males, despite being the dominant sex and showing higher rates of aggression. Drea [52] documented similar seasonal

patterns of androgen production in lemurs, with peak levels of androgens also occurring during the breeding season. In this study, both T and A_4 concentrations in female lemurs were lower than that of males (larger sex differences were apparent in T than in A_4). There were also major sex differences in the ratios of these two steroid hormones. T levels in males were twofold higher than A_4 but in females A_4 levels were approximately tenfold higher than T.

In lemurs, then, there is clear evidence of a temporal correspondence between elevated androgens during the breeding season and increased aggressive behaviour in females. In addition, A_4 /T ratios are proportionately higher in female lemurs than in males, suggesting that, like hyaenas, A_4 may serve as an important prehormone for androgenic effects on female aggression and dominance. However, the absence of a strong correlation between variation in levels of T and individual differences in female aggression and dominance is not consistent with the strong support for an exclusively activational role for T in lemurs.

4. Callitrichine primates (marmosets and tamarins)

Marmosets and tamarins offer a useful model for evaluating organizational and activational effects of steroid hormones on female aggressive behaviour. Callitrichine primates show a socially monogamous structure, including close heterosexual bonds [55]. Callitrichine social systems are also characterized by low rates of intragroup aggression, yet adult females are typically dominant over adult males using standard food competition paradigms [56]. Encounters between different groups, both in captivity and in the wild, are often characterized by aggressive territorial behaviour, especially between same-sex conspecifics [57], in which adult females play an active role. Behavioural responses of resident females to other female intruders can include physical attacks, but non-contact aggressive and threat displays by females are also common across species of this group ([57], see review in [58]). Thus, marmosets conform to our criteria for this paper: high levels of normative female aggression (in this case, in intergroup encounters) and dominance of females over males in adults.

(a) Prenatal androgen exposure

As marmosets and tamarins most often give birth to fraternal twins, initial speculation was that females would be less sensitive to androgens during gestation as they would run the risk of being overly masculinized by androgens secreted by their male womb-mates in the 50% of twin litters that were mixed sex. The impact of having a male versus female co-twin on female behavioural development has not been systematically examined to date. However, it is clear that marmoset fetuses are exposed to high and varying concentrations of maternally derived androgens *in utero* [31], with significant effects on later development. We demonstrated that differences in gestational exposure of fetuses to androgens influenced both morphological and behavioural development. First, offspring born to mothers with high gestational androgen levels in the first trimester were smaller at birth and had accelerated postnatal, especially juvenile, growth patterns [59]. In contrast to the results from the spotted hyaena that higher cub aggression and play-mounting was associated with high gestational

androgen exposure [17], marmosets offspring born to mothers with higher levels of maternal urinary androgens during late gestation displayed *decreased* rough-and-tumble play as juveniles [60]. While the effect was greater in males, females also showed reduced play behaviour as a consequence of high gestational exposure to androgens. These findings suggest that prenatal exposure to androgens affects at least two components of phenotypic characteristics in female marmosets, including rough-and-tumble play. Further, it is clear that sensitive periods for androgen-mediated changes in offspring morphological and behavioural phenotypes occur at different stages of gestation (growth: early gestation; behaviour: late gestation).

(b) Postnatal androgens

A single study has addressed the impact of immediate postnatal androgen exposure on later adult behavioural phenotypes in marmosets. Female common marmosets treated with T for the first 50 days of postnatal life showed higher rates of rough-and-tumble play during the juvenile period if they had a female co-twin (but not a male co-twin), suggesting that neonatally administered androgens may masculinize juvenile female behaviour. However, differences in the effects of androgen treatment as a function of the sex of the potential play partner indicate that there are social influences that modulate the behavioural consequences of elevated postnatal androgens in females [61,62]. These neonatal androgen manipulations also impacted sex-typical sexual behaviour. When neonatally T-treated females were tested as adults, they showed higher rates of male-typical courtship and copulatory behaviour with a receptive female marmoset than untreated females. Postnatal steroid environments also impact aggressive behaviour in tamarins. Female tamarins ovariectomized (OVX) prior to puberty exhibited fewer threat displays in an intruder paradigm test than intact females [63]. However, OVX and intact females showed no differences in physically injurious agonistic behaviour during intruder trials with unfamiliar females.

(c) Adult hormones and female aggressive behaviour

To our knowledge, only one study has addressed activational effects of androgens on aggression in adult female callitrichine primates. In Wied's black tufted-ear marmosets, baseline levels of androgens in adult females did not predict levels of aggression during an intruder trial with an unfamiliar female [64]. However, post-aggression levels of androgen measured at 6–24 h post-encounter were altered by the quality of the aggressive interaction by resident females. Those resident females that displayed high levels of aggression toward female intruders showed significant elevations in post-encounter urinary T, whereas females that showed low or no aggression toward the intruder had unaltered T levels. This aggression-dependent elevation in T is similar to the response of resident male marmosets to intruders in the post-encounter period [65] and suggests a facilitative or 'challenge-like' response [66] of androgen to high levels of aggression in both adult male and female marmosets.

5. Rodents

(a) Naked mole-rat (*Heterocephalus glaber*)

The mammalian species with perhaps the most dramatic system of female dominance and aggression is the naked

mole-rat. This subterranean eusocial mammal lives in colonies of up to 300 individuals, and breeding is typically monopolized by a single female (queen) and from one to three males [67]. Aggression and dominance is expressed primarily by intraburrow 'shoving', head-to-head contests of strength that results in the subordinate being displaced backward in the colony tunnels. Queens perform the vast majority of shoving (and never receive shoving from any colony member), breeding males show substantially lower rates and non-breeders of both sexes rarely engage in shoving. The queen's death or removal is associated with increased shoving and the rapid onset of ovarian function by the successor queen [68]. Consistent with an activational hypothesis, during queen transition periods, urinary T levels are threefold higher in aggressive females than in non-aggressive females [69]. Observations from a captive colony in which several non-breeding females engaged in shoving further supports the activational hypothesis, because urinary T levels among these females were positively correlated with aggressive shoving [69]. Areas of the brain that are typically sexually dimorphic in mammals (ventromedial nucleus of the hypothalamus, paraventricular nucleus and medial amygdala) are similar in size in breeding male and female naked mole-rats, and are larger in breeders of both sexes than in non-breeders [70]. Unfortunately, no data are available on pre- and postnatal hormone concentrations in this species, but the lability of ovarian function and subsequent steroid hormone *milieu* as a consequence of social status [68] suggests significant postpubertal plasticity in these otherwise sexually dimorphic brain regions.

(b) Cricetidae

Females of several seasonally breeding rodent species of the family Cricetidae display conspecific social aggression (in addition to maternal aggression; see Bosch [71]) as the result of organizational programming of androgens. Typically, female rodents exhibit significantly lower levels of aggression and androgens than males do. In typically male-dominant rodent species (e.g. brown rat), there are numerous pharmacological studies to suggest that gonadal steroid exposure plays a role in female aggressiveness [72,73]. Treatment with T and E₂ together contributes to the greatest increases in aggressiveness [74]. However, this treatment regimen is not sufficient to increase aggressiveness in typically non-aggressive female rats [75]. This suggests that a highly aggressive phenotype is influenced by gonadal steroids in male-dominant rodent species, but that females with phenotypically low aggression do not appear to be as strongly influenced by gonadal steroid metabolism. In rodent species where females naturally exhibit aggressive behaviour (i.e. Syrian hamsters and California mice), the relative influence of gonadal steroids on aggressive behaviour is dissimilar to gonadal steroid–aggression relationships in females of male-dominant rodent species.

(c) Syrian hamster (*Mesocricetus auratus*)

Female Syrian hamsters not only display significantly more aggressive behaviour than males [76], but also tend to dominate males in heterosexual encounters [77]. This sex difference may be attributable to relative differences in steroid metabolism between the sexes [76]. Neonatal androgen and oestrogen exposure, as well as variation in steroid and neuropeptide levels in adulthood, have been implicated in the expression of female intrasexual social aggression. Gonadal

steroid exposure during development can have a modulatory effect on behavioural phenotypes in females later in life. Female Syrian hamsters treated with T on the second day post-birth showed increased aggressiveness relative to control females [78]. This suggests that T exposure for females during development leads to a higher expression of aggressive behaviour later in life, thus providing support for an organizational hypothesis.

Additionally, changes in oestrous state have pronounced effects on the expression of aggressive behaviour in Syrian hamsters, suggesting that ovarian hormones can alter female aggressiveness. Non-oestrous females display high levels of conspecific aggression, while aggressive behaviour is essentially absent in oestrous females [79]. These fluctuations in aggressive behaviour across the oestrus cycle are attributable to changes in oestrogen and progesterone (P₄). In females treated with both E₂ and P₄, aggression is completely suppressed [79,80]. However, this reduction in aggressive behaviour is only seen when E₂ and P₄ levels are increased in concert, and is not seen when these hormones are replaced individually [79,80]. The relative contributions of gonadal steroids in these atypical rodent species appear to influence female aggression differently than in more typical male-dominant rodents. However, aggressive encounters between unisexual pairs of female Syrian hamsters were unaffected by gonadectomy, and treatment with T [81] and oestrogen [82]. Thus, despite the evidence suggesting that female aggression is influenced by hormonal state, the aggression-inducing effect of gonadal steroids on females does not appear to apply across all contexts.

(d) California mouse (*Peromyscus californicus*)

Female California mice show a similar aggression phenotype as males, with increased aggression during winter-like short days [83]. There are important consequences of androgen exposure on the development of neural mechanisms for aggression [84]. Female mice that were exposed to T neonatally, and then treated with T in adulthood displayed increased levels of aggression, relative to females that did not receive neonatal treatment [84]. Furthermore, neonatal treatment with oestrogen also greatly increased the expression of female aggression in adulthood, mirroring the effect seen from neonatal T treatment [85]. Neonatal females treated with T during development appear to have their neural system primed for aggression, by increasing androgen receptor sensitivity to T in adulthood [84]. This increased sensitivity to T leads to higher levels of female aggression displayed during adulthood.

There is also substantial evidence that indicates androgens have an activational effect on the expression of aggression in female rodents. In female California mice, T facilitates baseline aggression, as expected, but E₂ inhibits aggressive behaviour, as oestradiol treatment given to ovariectomized female mice did not induce aggressive behaviour [86]. Thus, it may be that the E₂/T ratio is more important than either hormone alone as an endocrine modulating factor for baseline aggression in this species [87]. Furthermore, female California mice displayed a decrease in P₄ levels, and the P₄/T ratio, following an intruder encounter, suggesting that P₄ inhibits female intrasexual aggression. It was also suggested that a post-encounter decrease in the P₄/T ratio may serve to modify aggression in future encounters [87].

Recent work has found that the underlying neuroendocrine mechanism responsible for increased aggression

owing to T treatment is the downregulation of neurosteroid biosynthesis of allopregnanolone (AP). Predictably, female Swiss Webster mice that were administered T displayed an increase in the expression of aggression. However, this long-term treatment with T also led to a downregulation of brain AP levels [88]. Interestingly, this T-induced aggression can be reversed by selectively increasing AP content [88] suggesting that endogenous pregnane steroids play an important modulatory role in attenuation of female aggression. Attention to androgen modulation of neurosteroid synthesis and function might provide additional insights into mechanisms of agonistic behaviour in mammals with sex-reversed patterns of aggression.

6. Rock hyrax (*Procavia capensis*)

Rock hyrax social structure is characterized by multi-male–multi-female groups, with a small number of females ‘monopolized’ for reproduction by a territorial male. Females do engage in cooperative rearing of offspring, although relatedness among cooperating females is not known. Some aggression within hyrax groups occurs in intrasexual contexts, but the majority of agonistic interactions, including threat displays, chases and avoidance, occurs between males (both territorial and non-group bachelor males) and females. In spite of the reproductive ‘monopolization’ by territorial males, adult female rock hyrax in most wild groups under observation are behaviourally dominant to males [89].

Across populations of rock hyrax, levels of T in females (measured in hair samples) are similar to or higher than levels in males [89,90] suggesting that T may serve an activational role in female aggression. However, while T was a strong predictor of within-group differences in male dominance status (and body size), individual differences in female aggression rates and dominance were independent of T levels. This finding is not consistent with the potential activational link between androgens and female aggression. Accounting for high T concentrations in female rock hyrax poses a dilemma, because there appear to be significant costs for females without apparent benefits. Females with elevated androgens have a larger proportion of skin absent of hair, a proxy measure for ectoparasite load and/or susceptibility to infection, yet other than female dominance over males, individual differences in T are not correlated with enhanced rates of aggression or reproductive success in females, as measured by litter size [90].

7. Summary, conclusion and future directions

This review has highlighted the role of androgens in shaping mammalian female aggressive patterns and female dominance in a cluster of species that depart from typical male bias in both these measures. An overall summary of the empirical data presented in this paper is highlighted in table 1. It is clear from this review that androgenic steroids play an important, but not exclusive, role in shaping these phenotypes. It is equally clear that our understanding of the role of these steroid hormones in both organizing and activating female aggression is far from complete. None of the species listed in this review have the breadth and depth of research on these issues that characterizes more traditional mammals, especially those that are studied extensively in the laboratory or in captive situations (e.g. rodents: [93]; macaque monkeys: [94]).

Table 1 Summary of female aggression, dominance, and organizational and activational effects of androgen hormones in ‘atypical’ mammals.

species	aggression	dominance	organizational effect	activational effects	facilitative effects	T	A ₄	comments	references
hyena	F > M	F > M	+	n.a.	n.a.	M > F	F > M		[17,18,21,24]
lemur	F > M	F > M	+	–	n.a.	M > F	M ≅ F	mixed results for activation of aggression and dominance	[49,50,52,53]
marmoset	M ≅ F	F > M	+	–	+	M > F	n.a.		[60–64]
naked mole-rat	F > M	F > M	n.a.	+	n.a.	M > F	n.a.	dominant female appears to suppress subordinate reproductive hormones	[68,69]
Syrian hamster	F > M	F > M	+	–	n.a.	M > F	n.a.	mixed results for gonadal steroid exposure effect on aggression. Photoperiod important as well	[78,81,82]
California mouse	M ≅ F	n.a.	n.a.	+	+	M > F	n.a.	E ₂ /T ratio more important than either hormone alone as contributors influencing baseline aggression	[87,91,92]
rock hyrax	F > M	F > M	n.a.	–	n.a.	F > M	n.a.		[89,90]

The potential for significant peripubertal organizational effects of androgens on female aggression, as highlighted in the work of Sisk and co-workers [95], in more conventional mammalian species has yet to be explored. Further, the potential for androgens to play a facilitative role in future aggression in females as a consequence of aggression-induced rises in these hormones [66] has only been addressed in a limited number of these species. Additional species that depart from mammalian norms regarding sexual dimorphism in aggression, although not to the extent to which our focal species do, are also worthy of further study. For instance, unlike male-dominant and -aggressive chimpanzees (*Pan troglodytes*), male and female roles in bonobos (*Pan paniscus*) are relatively egalitarian, and females tend to be dominant over males, although without the overt aggression associated with species like spotted hyaenas [96]. Comparative analyses of androgen levels in adults show that male : female ratios of urinary androgens are highly male-biased in chimpanzees, but more similar between the sexes in bonobos [97]. However, these differences appear to be attributable primarily to elevated T in male chimpanzees, rather than elevated T in female bonobos. A recent analysis of digit ratios (2D : 4D; a proxy for *in utero* exposure to androgens) in chimpanzees and bonobos failed to find evidence for elevated prenatal exposure to androgens in female bonobos [98], shedding doubt on an organizational effect on the propensity toward female dominance in this species.

However, further work is required to clarify both activational and organizational effects of androgens in bonobos and other species with less dramatic but still female-biased dominance and aggression.

Finally, more sophisticated analyses of endocrine effects on brain structure underlying agonistic behaviour [26,70], increased knowledge of the enzymatic substrates that mediate important transitions from less active to more active steroid moieties (and vice versa; [15,28]) and an appreciation that variation in carrier peptides is critical for estimating the actual exposure of steroid-sensitive brain mechanisms to hormones [36] will enrich our understanding of gonadal influences on aggressive behaviour. What is perfectly clear, however, is that the kind of variation in mechanism-behaviour relationships revealed in this review, which has its roots in the operation of natural selection on these systems, can provide us with knowledge as rich and informative as data gleaned from induced genetic variation from steroid receptor knock-out models and other laboratory constructs [99,100].

Acknowledgements. We thank the guest editors for the invitation to participate in this Special Issue on Female Aggression, and anonymous reviewers for useful and constructive comments on previous versions of the paper.

Funding statement. The Callitrichid Research Center at the University of Nebraska at Omaha is supported in part by funds from the National Institutes of Health (HD042882) and the Nebraska Research Initiative.

References

- Darwin CH. 1871 *The descent of man, and selection in relation to sex*. Princeton, NY: Princeton University Press.
- Andersson MB. 1994 *Sexual selection*. Princeton, NY: Princeton University Press.
- Trivers R. 1972 Parental investment and sexual selection. In *Sexual selection and the descent of man 1871–1971* (ed. B Campbell), pp. 137–179. Chicago, IL: Aldine.
- Karubian J, Swaddle JP. 2001 Selection on females can create 'larger males'. *Proc. R. Soc. Lond. B* **268**, 725–728. (doi:10.1098/rspb.2000.1407)
- Loison A, Gaillard J-M, Pelabon C, Yoccoz NG. 1999 What factors shape sexual size dimorphism in ungulates? *Evol. Ecol. Res.* **1**, 611–633.
- Ralls K. 1976 Mammals in which females are larger than males. *Q. Rev. Biol.* **51**, 245–276. (doi:10.1086/409310)
- Clutton-Brock T. 2009 Sexual selection in females. *Anim. Behav.* **77**, 3–11. (doi:10.1016/j.anbehav.2008.08.026)
- Isaac JL. 2005 Potential causes and life-history consequences of sexual size dimorphism in mammals. *Mammal Rev.* **35**, 101–115. (doi:10.1111/j.1365-2907.2005.00045.x)
- Clutton-Brock TH, Huchard E. 2013 Social competition and selection in males and females. *Phil. Trans. R. Soc. B* **368**, 20130074. (doi:10.1098/rstb.2013.0074)
- Wallen K. 2005 Hormonal influences on sexually differentiated behavior in nonhuman primates. *Front. Neuroendocrinol.* **26**, 7–26. (doi:10.1016/j.yfrne.2005.02.001)
- Hau M. 2007 Regulation of male traits by testosterone: implications for the evolution of vertebrate life histories. *BioEssays* **29**, 133–144. (doi:10.1002/bies.20524)
- Soma KK, Scotti M-AL, Newman AEM, Charlier TD, Demas GE. 2008 Novel mechanisms for neuroendocrine regulation of aggression. *Front. Neuroendocrinol.* **29**, 476–489. (doi:10.1016/j.yfrne.2007.12.003)
- Glickman SE, Cunha GR, Drea CM, Conley AJ, Place NJ. 2006 Mammalian sexual differentiation: lessons from the spotted hyena. *Trends Endocrinol. Metab.* **17**, 349–356. (doi:10.1016/j.tem.2006.09.005)
- Glickman SE, Frank LG, Davidson JM, Smith ER, Siiteri PK. 1987 Androstenedione may organize or activate sex-reversed traits in female spotted hyenas. *Proc. Natl. Acad. Sci. USA* **84**, 3444–3447. (doi:10.1073/pnas.84.10.3444)
- Yalcinkaya TM, Siiteri PK, Vigne J-L, Licht P, Pavgi S, Frank LG, Glickman SE. 1993 A mechanism for virilization of female spotted hyenas *in utero*. *Science* **260**, 1929–1931. (doi:10.1126/science.8391165)
- Dloniak SM, French JA, Place NJ, Weldele ML, Glickman SE, Holekamp KE. 2004 Non-invasive monitoring of fecal androgens in spotted hyenas (*Crocuta crocuta*). *Gen. Comp. Endocrinol.* **135**, 51–61. (doi:10.1016/j.ygcen.2003.08.011)
- Dloniak SM, French JA, Holekamp KE. 2006 Rank-related maternal effects of androgens on behaviour in wild spotted hyenas. *Nature* **440**, 1190–1193. (doi:10.1038/nature04540)
- Holekamp KE, Swanson EM, Van Meter PE. 2013 Developmental constraints on behavioural flexibility. *Phil. Trans. R. Soc. B* **368**, 20120350. (doi:10.1098/rstb.2012.0350)
- Place NJ, Holekamp KE, Sisk CL, Weldele ML, Coscia EM, Drea CM, Glickman SE. 2002 Effects of treatment with anti-androgens on LH secretion and gonadal steroidogenesis in the spotted hyena *Crocuta crocuta*. *Biol. Reprod.* **67**, 1405–1413. (doi:10.1095/biolreprod.102.004226)
- Drea CM. 2009 Endocrine mediators of masculinization in female mammals. *Curr. Dir. Psychol. Sci.* **18**, 221–226. (doi:10.1111/j.1467-8721.2009.01640.x)
- Frank LG, Glickman SE, Licht P. 1991 Fatal sibling aggression, precocial development, and androgens in neonatal spotted hyenas. *Science* **252**, 702–704. (doi:10.1126/science.2024122)
- Golla W, Hofer H, East ML. 1999 Within-litter sibling aggression in spotted hyenas: effect of maternal nursing, sex and age. *Anim. Behav.* **58**, 715–726. (doi:10.1006/anbe.1999.1189)
- Glickman SE, Frank LG, Pavgi S, Licht P. 1992 Hormonal correlates of 'masculinization' in female spotted hyenas (*Crocuta crocuta*). 1. Infancy to sexual maturity. *J. Reprod. Fertil.* **95**, 451–462. (doi:10.1530/jrf.0.0950451)
- Goymann W, East ML, Hofer H. 2001 Androgens and the role of female 'hyperaggressiveness' in spotted hyenas (*Crocuta crocuta*). *Horm. Behav.* **39**, 83–92. (doi:10.1006/hbeh.2000.1634)

25. Fenstemaker SB, Zup SL, Frank LG, Glickman SE, Forger NG. 1999 A sex difference in the hypothalamus of the spotted hyena. *Nat. Neurosci.* **2**, 943–945. (doi:10.1038/14728)
26. Rosen GJ, De Vries GJ, Villalba C, Weldele ML, Place NJ, Coscia EM, Glickman SE, Forger NG. 2006 Distribution of vasopressin in the forebrain of spotted hyenas. *J. Comp. Neurol.* **498**, 80–92. (doi:10.1002/cne.21032)
27. Dloniak SM, French JA, Holekamp KE. 2006 Faecal androgen concentrations in adult male spotted hyaenas, *Crocuta crocuta*, reflect interactions with socially dominant females. *Anim. Behav.* **71**, 27–37. (doi:10.1016/j.anbehav.2005.03.016)
28. Browne P, Place NJ, Vidal JD, Moore IT, Cunha GR, Glickman SE, Conley AJ. 2006 Endocrine differentiation of fetal ovaries and testes of the spotted hyena (*Crocuta crocuta*): timing of androgen-independent versus androgen-driven genital development. *Reproduction* **132**, 649–659. (doi:10.1530/rep.1.01120)
29. Gleicher N, Weghofer A, Barad DH. 2011 The role of androgens in follicle maturation and ovulation induction: friend or foe of infertility treatment. *Reprod. Biol. Endocrinol.* **9**, 116. (doi:10.1186/1477-7827-9-116)
30. Fite JE, French JA, Patera KJ, Hopkins EC, Rukstalis M, Ross CN. 2005 Elevated urinary testosterone excretion and decreased maternal caregiving effort in marmosets when conception occurs during the period of infant dependence. *Horm. Behav.* **47**, 39–48. (doi:10.1016/j.yhbeh.2004.08.005)
31. French JA, Smith AS, Birnie AK. 2010 Maternal gestational androgen levels in female marmosets (*Callithrix geoffroyi*) vary across trimesters but do not vary with the sex ratio of litters. *Gen. Comp. Endocrinol.* **165**, 309–314. (doi:10.1016/j.yggen.2009.07.015)
32. Carroll TB, Aron DC, Findling JW, Tyrrell JB. 2011 Glucocorticoids and adrenal androgens. In *Greenspan's basic & clinical endocrinology* (eds DG Gardner, D Shoback), pp. 285–327. New York, NY: McGraw-Hill Medical.
33. Busso JM, Ruiz RD. 2011 Excretion of steroid hormones in rodents: an overview on species differences for new biomedical animal research models. In *Contemporary aspects of endocrinology* (ed. E Diamanti-Kandarakis), pp. 375–396. Intech Open Access. (doi:10.5772/17051)
34. Pozo OJ, Marcos J, Ventura R, Fabregat A, Segura J. 2010 Testosterone metabolism revisited: discovery of new metabolites. *Anal. Bioanal. Chem.* **398**, 1759–1770. (doi:10.1007/s00216-010-4082-0)
35. Hammond GL. 2011 Diverse roles for sex hormone-binding globulin in reproduction. *Biol. Reprod.* **85**, 431–441. (doi:10.1095/biolreprod.111.092593)
36. Hammond GL, Miguel-Queralto S, Yalcinkaya TM, Underhill C, Place NJ, Glickman SE, Drea CM, Wagner AP, Siiteri PK. 2012 Phylogenetic comparisons implicate sex hormone-binding globulin in 'masculinization' of the female spotted hyena (*Crocuta crocuta*). *Endocrinology* **153**, 1435–1443. (doi:10.1210/en.2011-1837)
37. Jolly A. 1966 Lemur social behavior and primate intelligence. *Science* **153**, 501–506. (doi:10.1126/science.153.3735.501)
38. Kappeler PM. 1993 Reconciliation and post-conflict behaviour in ringtailed lemurs, *Lemur catta* and redfronted lemurs, *Eulemur fulvus rufus*. *Anim. Behav.* **45**, 901–915. (doi:10.1006/anbe.1993.1110)
39. Jolly A. 1998 Pair-bonding, female aggression and the evolution of lemur societies. *Folia Primatol. (Basel)* **69**, 1–13. (doi:10.1159/000052693)
40. Pollock J. 1979 Spatial distribution and ranging behaviour in lemurs. In *The Study of Prosimian Behavior* (eds GA Doyle, RD Martin), pp. 359–409. London, UK: Academic Press.
41. Kappeler PM. 1993 Variation in social structure: the effects of sex and kinship on social interactions in three lemur species. *Ethology* **93**, 125–145. (doi:10.1111/j.1439-0310.1993.tb00984.x)
42. Pereira ME, Kappeler PM. 1997 Divergent systems of agonistic behaviour in lemurid primates. *Behaviour* **134**, 225–274. (doi:10.1163/156853997X00467)
43. Pereira ME. 1995 Development and social dominance among group-living primates. *Am. J. Primatol.* **37**, 143–175. (doi:10.1002/ajp.1350370207)
44. Mitani JC, Call J, Kappeler PM, Palombit RA, Silk JB (eds). 2012 *The evolution of primate societies*. Chicago, IL: University of Chicago Press.
45. Vick LG, Pereira ME. 1989 Episodic targeting aggression and the histories of lemur social groups. *Behav. Ecol. Sociobiol.* **25**, 3–12. (doi:10.1007/BF00299705)
46. Digby L, Mclean Stevens A. 2007 Maintenance of female dominance in blue-eyed black lemurs (*Eulemur macaco flavifrons*) and gray bamboo lemurs (*Hapalemur griseus griseus*) under semi-free-ranging and captive conditions. *Zoo Biol.* **26**, 345–361. (doi:10.1002/zoo.20140)
47. White FJ, Overdorff DJ, Keith-Lucas T, Rasmussen MA, Eddie Kallam W, Forward Z. 2007 Female dominance and feeding priority in a prosimian primate: experimental manipulation of feeding competition. *Am. J. Primatol.* **69**, 295–304. (doi:10.1002/ajp.20346)
48. Drea CM, Weil A. 2008 External genital morphology of the ring-tailed lemur (*Lemur catta*): females are naturally 'masculinized'. *J. Morphol.* **269**, 451–463. (doi:10.1002/jmor.10594)
49. Drea CM. 2011 Endocrine correlates of pregnancy in the ring-tailed lemur (*Lemur catta*): implications for the masculinization of daughters. *Horm. Behav.* **59**, 417–427. (doi:10.1016/j.yhbeh.2010.09.011)
50. Ostner J, Heistermann M, Kappeler PM. 2003 Intersexual dominance, masculinized genitals and prenatal steroids: comparative data from lemurid primates. *Naturwissenschaften* **90**, 141–144.
51. Monaghan EP, Glickman SE. 1992 Hormones and aggressive behavior. *Behav. Endocrinol.* 261–285.
52. Drea CM. 2007 Sex and seasonal differences in aggression and steroid secretion in *Lemur catta*: are socially dominant females hormonally 'masculinized'? *Horm. Behav.* **51**, 555–567. (doi:10.1016/j.yhbeh.2007.02.006)
53. Von Engelhard N, Kappeler PM, Heistermann M. 2000 Androgen levels and female social dominance in *Lemur catta*. *Proc. R. Soc. Lond. B* **267**, 1533–1539. (doi:10.1098/rspb.2000.1175)
54. Cavigelli SA, Pereira ME. 2000 Mating season aggression and fecal testosterone levels in male ring-tailed lemurs (*Lemur catta*). *Horm. Behav.* **37**, 246–255. (doi:10.1006/hbeh.2000.1585)
55. Agmo A, Smith AS, Birnie AK, French JA. 2012 Behavioral characteristics of pair bonding in the black tufted-ear marmoset (*Callithrix penicillata*). *Behaviour* **149**, 407. (doi:10.1163/156853912X638454)
56. Tardif S, Richter C. 1981 Competition for a desired food in family groups of the common marmoset (*Callithrix jacchus*) and the cotton-top tamarin (*Saguinus oedipus*). *Lab. Anim. Sci.* **31**, 52–55.
57. Lazaro-Perea C. 2001 Intergroup interactions in wild common marmosets, *Callithrix jacchus*: territorial defence and assessment of neighbours. *Anim. Behav.* **62**, 11–21. (doi:10.1006/anbe.2000.1726)
58. French JA, Inglett BJ. 1991 Responses to novel social stimuli in callitrichid monkeys: a comparative perspective. In *Primate responses to environmental change* (ed. HO Box), pp. 275–294. New York, NY: Chapman and Hall.
59. Smith AS, Birnie AK, French JA. 2010 Maternal androgen levels during pregnancy are associated with early-life growth in Geoffroy's marmosets, *Callithrix geoffroyi*. *Gen. Comp. Endocrinol.* **166**, 307–313. (doi:10.1016/j.yggen.2009.10.008)
60. Birnie AK, Hendricks SE, Smith AS, Milam R, French JA. 2012 Maternal gestational androgens are associated with decreased juvenile play in white-faced marmosets (*Callithrix geoffroyi*). *Horm. Behav.* **62**, 136–145. (doi:10.1016/j.yhbeh.2012.05.014)
61. Abbott DH, Hearn JP. 1978 The effects of neonatal T on female monkeys. *Ciba Found. Symp.* **62**, 299–327.
62. Abbott DH. 1984 Differentiation of sexual behavior in female marmoset monkeys: effects of neonatal testosterone or a male co-twin. *Prog. Brain Res.* **61**, 349–358. (doi:10.1016/S0079-6123(08)64446-5)
63. Epple G. 1982 Effects of prepubertal ovariectomy on the development of scent glands, scent marking, and aggressive behaviors of female tamarin monkeys (*Saguinus fuscicollis*). *Horm. Behav.* **16**, 330–342. (doi:10.1016/0018-506X(82)90031-9)
64. Ross CN, French JA. 2011 Female marmosets' behavioral and hormonal responses to unfamiliar intruders. *Am. J. Primatol.* **73**, 1072–1081.
65. Ross CN, French JA, Patera KJ. 2004 Intensity of aggressive interactions modulates testosterone in male marmosets. *Physiol. Behav.* **83**, 437–445. (doi:10.1016/j.physbeh.2004.08.036)
66. Wingfield JC, Hegner RE, Dufty Jr AM, Ball GF. 1990 The 'challenge hypothesis': theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *Am. Nat.* **136**, 829–846. (doi:10.1086/285134)
67. Jarvis JU. 1981 Eusociality in a mammal: cooperative breeding in naked mole-rat colonies.

- Science* **212**, 571–573. (doi:10.1126/science.7209555)
68. Margulis SW, Saltzman W, Abbott DH. 1995 Behavioral and hormonal changes in female naked mole-rats (*Heterocephalus glaber*) following removal of the breeding female from a colony. *Horm. Behav.* **29**, 227–247. (doi:10.1006/hbeh.1995.1017)
 69. Clarke FM, Faulkes CG. 1997 Dominance and queen succession in captive colonies of the eusocial naked mole-rat, *Heterocephalus glaber*. *Proc. R. Soc. Lond. B* **264**, 993–1000. (doi:10.1098/rspb.1997.0137)
 70. Holmes MM, Goldman BD, Forger NG. 2008 Social status and sex independently influence androgen receptor expression in the eusocial naked mole-rat brain. *Horm. Behav.* **54**, 278–285. (doi:10.1016/j.yhbeh.2008.03.010)
 71. Bosch OJ. 2013 Maternal aggression in rodents: brain oxytocin and vasopressin mediate pup defence. *Phil. Trans. R. Soc. B* **368**, 20130085. (doi:10.1098/rstb.2013.0085)
 72. Albert DJ, Jonik RH, Walsh ML. 1990 Aggression by ovariectomized female rats with testosterone implants: competitive experience activates aggression toward unfamiliar females. *Physiol. Behav.* **47**, 699–703. (doi:10.1016/0031-9384(90)90080-N)
 73. Albert DJ, Jonik RH, Walsh ML. 1990 Hormone-dependent aggression in female rats: testosterone implants attenuate the decline in aggression following ovariectomy. *Physiol. Behav.* **47**, 659–664. (doi:10.1016/0031-9384(90)90074-E)
 74. Albert DJ, Jonik RH, Walsh ML. 1991 Hormone-dependent aggression in the female rat: testosterone plus estradiol implants prevent the decline in aggression following ovariectomy. *Physiol. Behav.* **49**, 673–677. (doi:10.1016/0031-9384(91)90300-D)
 75. Albert DJ, Jonik RH, Walsh ML. 1993 Influence of combined estradiol and testosterone implants on the aggressiveness of nonaggressive female rats. *Physiol. Behav.* **53**, 709–713. (doi:10.1016/0031-9384(93)90177-H)
 76. Taravosh-Lahn K, Delville Y. 2004 Aggressive behavior in female golden hamsters: development and the effect of repeated social stress. *Horm. Behav.* **46**, 428–435. (doi:10.1016/j.yhbeh.2004.03.007)
 77. Payne AP, Swanson HH. 1970 Agonistic behavior between pairs of hamsters of the same and opposite sex in a neutral observation area. *Behaviour* **36**, 259–267. (doi:10.1163/156853970X00402)
 78. Payne AP. 1976 A comparison of the effects of neonatally administered testosterone, testosterone propionate and dihydrotestosterone on aggressive and sexual behaviour in the female golden hamster. *J. Endocrinol.* **69**, 23–31. (doi:10.1677/joe.0.0690023)
 79. Floody OR, Pfaff DW. 1977 Aggressive behavior in female hamsters: the hormonal basis for fluctuations in female aggressiveness correlated with estrous state. *J. Comp. Physiol. Psychol.* **91**, 443–464. (doi:10.1037/h0077341)
 80. Meisel RL, Sterner MR, Diekman MA. 1988 Differential hormonal control of aggression and sexual behavior in female Syrian hamsters. *Horm. Behav.* **22**, 453–466. (doi:10.1016/0018-506X(88)90050-5)
 81. Payne AP, Swanson HH. 1972 The effect of sex hormones on the aggressive behaviour of the female golden hamster (*Mesocricetus auratus* Waterhouse). *Anim. Behav.* **20**, 782–787. (doi:10.1016/S0003-3472(72)80152-0)
 82. Vandenberg JG. 1971 The effects of gonadal hormones on the aggressive behaviour of adult golden hamsters (*Mesocricetus auratus*). *Anim. Behav.* **19**, 589–594. (doi:10.1016/S0003-3472(71)80116-1)
 83. Silva AL, Fry WHD, Sweeney C, Trainor BC. 2010 Effects of photoperiod and experience on aggressive behavior in female California mice. *Behav. Brain Res.* **208**, 528–534. (doi:10.1016/j.bbr.2009.12.038)
 84. Edwards DA. 1969 Early androgen stimulation and aggressive behavior in male and female mice. *Physiol. Behav.* **4**, 333–338. (doi:10.1016/0031-9384(69)90185-1)
 85. Edwards DA, Herndon J. 1970 Neonatal estrogen stimulation and aggressive behavior in female mice. *Physiol. Behav.* **5**, 993–995. (doi:10.1016/0031-9384(70)90153-8)
 86. Edwards DA, Burge KG. 1971 Estrogenic arousal of aggressive behavior and masculine sexual behavior in male and female mice. *Horm. Behav.* **2**, 239–245. (doi:10.1016/0018-506X(71)90021-3)
 87. Davis ES, Marler CA. 2003 The progesterone challenge: steroid hormone changes following a simulated territorial intrusion in female *Peromyscus californicus*. *Horm. Behav.* **44**, 185–198. (doi:10.1016/S0018-506X(03)00128-4)
 88. Pinna G, Costa E, Guidotti A. 2005 Changes in brain testosterone and allopregnanolone biosynthesis elicit aggressive behavior. *Proc. Natl Acad. Sci. USA* **102**, 2135–2140. (doi:10.1073/pnas.0409643102)
 89. Koren L, Mokady O, Geffen E. 2006 Elevated testosterone levels and social ranks in female rock hyrax. *Horm. Behav.* **49**, 470–477. (doi:10.1016/j.yhbeh.2005.10.004)
 90. Koren L, Geffen E. 2009 Androgens and social status in female rock hyraxes. *Anim. Behav.* **77**, 233–238. (doi:10.1016/j.anbehav.2008.09.031)
 91. Barkley MS, Goldman BD. 1977 Testosterone-induced aggression in adult female mice. *Horm. Behav.* **9**, 76–84. (doi:10.1016/0018-506X(77)90052-6)
 92. Perch e F, Young J, Robel P, Simon NG, Haug M. 2001 Prenatal testosterone treatment potentiates the aggression-inhibiting effect of the neurosteroid dehydroepiandrosterone in female mice. *Aggress. Behav.* **27**, 130–138. (doi:10.1002/ab.5)
 93. Lenz KM, Nugent BM, McCarthy MM. 2012 Sexual differentiation of the rodent brain: dogma and beyond. *Front. Neurosci.* **6**, 26. (doi:10.3389/fnins.2012.00026)
 94. Thornton J, Zehr JL, Loose MD. 2009 Effects of prenatal androgens on rhesus monkeys: a model system to explore the organizational hypothesis in primates. *Horm. Behav.* **55**, 633–644. (doi:10.1016/j.yhbeh.2009.03.015)
 95. Schulz KM, Molenda-Figueira HA, Sisk CL. 2009 Back to the future: the organizational-activational hypothesis adapted to puberty and adolescence. *Horm. Behav.* **55**, 597–604. (doi:10.1016/j.yhbeh.2009.03.010)
 96. De Waal FB, Lanting F. 1997 *Bonobo: the forgotten ape*. Berkeley, CA: University of California Press.
 97. Sannen A, Heistermann M, Van Elsacker L, M hle U, Eens M. 2003 Urinary testosterone metabolite levels in bonobos: a comparison with chimpanzees in relation to social system. *Behaviour* **140**, 683–696. (doi:10.1163/156853903322149504)
 98. McIntyre MH, Herrmann E, Wobber V, Halbax M, Mohamba C, de Sousa N, Atencia R, Cox D, Hare B. 2009 Bonobos have a more human-like second-to-fourth finger length ratio (2D:4D) than chimpanzees: a hypothesized indication of lower prenatal androgens. *J. Hum. Evol.* **56**, 361–365. (doi:10.1016/j.jhevol.2008.12.004)
 99. Nelson RJ, Chiavegatto S. 2000 Aggression in knockout mice. *Inst. Lab Anim. Res. J.* **41**, 153–162. (doi:10.1093/ilr.41.3.153)
 100. Honda S-I, Wakatsuki T, Harada N. 2011 Behavioral analysis of genetically modified mice indicates essential roles of neurosteroidal estrogen. *Front. Endocrinol.* **2**, 40. (doi:10.3389/fendo.2011.00040)