

Assembling the Tree of the Monocotyledons: Plastome Sequence Phylogeny and Evolution of Poales

Author(s) :Thomas J. Givnish, Mercedes Ames, Joel R. McNeal, Michael R. McKain, P. Roxanne Steele, Claude W. dePamphilis, Sean W. Graham, J. Chris Pires, Dennis W. Stevenson, Wendy B. Zomlefer, Barbara G. Briggs, Melvin R. Duvall, Michael J. Moore, J. Michael Heaney, Douglas E. Soltis, Pamela S. Soltis, Kevin Thiele, and James H. Leebens-Mack

Source: *Annals of the Missouri Botanical Garden*, 97(4):584-616. 2010.

Published By: Missouri Botanical Garden

DOI:

URL: <http://www.bioone.org/doi/full/10.3417/2010023>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

ASSEMBLING THE TREE OF THE MONOCOTYLEDONS: PLASTOME SEQUENCE PHYLOGENY AND EVOLUTION OF POALES¹

Thomas J. Givnish,² Mercedes Ames,² Joel R. McNeal,³ Michael R. McKain,³ P. Roxanne Steele,⁴ Claude W. dePamphilis,⁵ Sean W. Graham,⁶ J. Chris Pires,⁴ Dennis W. Stevenson,⁷ Wendy B. Zomlefer,³ Barbara G. Briggs,³ Melvin R. Duval,⁹ Michael J. Moore,¹⁰ J. Michael Heaney,¹¹ Douglas E. Soltis,¹¹ Pamela S. Soltis,¹² Kevin Thiele,¹³ and James H. Leebens-Mack³

ABSTRACT

The order Poales comprises a substantial portion of plant life (7% of all angiosperms and 33% of monocots) and includes taxa of enormous economic and ecological significance. Molecular and morphological studies over the past two decades, however, leave uncertain many relationships within Poales and among allied commelinid orders. Here we present the results of an initial project by the Monocot AToL (Angiosperm Tree of Life) team on phylogeny and evolution in Poales, using sequence data for 81 plastid genes (exceeding 101 aligned kb) from 83 species of angiosperms. We recovered highly concordant relationships using maximum likelihood (ML) and maximum parsimony (MP), with 98.2% mean ML bootstrap support across monocots. For the first time, ML resolves ties among Poales and other commelinid orders with moderate to strong support. Analyses provide strong support for Bromeliaceae being sister to the rest of Poales; Typhaceae, Rapateaceae, and cyperids (sedges, rushes, and their allies) emerge next along the phylogenetic spine. Graminids (grasses and their allies) and restiids (Restionaceae and its allies) are well supported as sister taxa. MP identifies a xyrid clade (Eriocaulaceae, Mayacaceae, Xyridaceae) sister to cyperids, but ML (with much stronger support) places them as a grade with respect to restiids + graminids. The conflict in resolution between these analyses likely reflects long-branch attraction and highly elevated substitution rates in some Poales. All other familial relationships within the order are strongly supported by both MP and ML analyses. Character-state mapping implies that ancestral Poales lived in sunny, fire-prone, at least seasonally damp/wet, and possibly nutrient-poor sites, and were animal pollinated. Five subsequent shifts to wind pollination—in Typhaceae, cyperids, restiids, Eceidaceae, and the vast PACCMAD-BEP clade of grasses—are significantly correlated with shifts to open habitats and small, inconspicuous, unisexual, and nectar-free flowers. Prime ecological movers driving the repeated evolution of wind pollination in Poales appear to include open habitats combined with the high local dominance of conspecific taxa, with the latter resulting from large-scale disturbances, combined with tall plant stature, vigorous vegetative

¹ This research was supported by National Science Foundation (NSF) grants DEB-0829762, DEB-0829849, DEB-0829868, DEB-0830009, DEB-0830020, and DEB-0830036 to members of the Monocot AToL Team, and by the Hertel Gift Fund of the University of Wisconsin-Madison to T.J.G. Jannice Friedman, Jana Vamosi, Spencer Barrett, and Kathleen Kay kindly provided access to data sets on wind pollination and related traits across angiosperms, and George Weiblen graciously let us use his data set on the occurrence of dioecy across monocots. Lynn Clark kindly shared information on bamboo seeds and pollination. Marco Duretto and Greg Jordan helped provide permits and suggest collecting locations in Tasmania; Joanne Birch provided assistance and camaraderie while collecting at Cradle Mountain. Godfrey Bourne and Margaret Chan-a-Sue were especially generous in helping to obtain research permits and organize transport to and from Kaieteur Falls National Park in Guyana. We would like to thank Frank Zich of the Australian Tropical Herbarium and Mark Clements of the Centre for Biodiversity Research for providing access to *Apostasia* Blume, and Jeremy Bruhl for sending material of *Neoastelia* J. B. Williams. Andrew Meade, Mark Pagel, Mark Chase, Ken Sytsma, Marie Adams, and Eric Cabot kindly provided technical assistance. Mike Havey generously provided access to important lab equipment. Chang Liu helped prepare DNA samples for sequencing. Raj Ayyampalayam wrote Perl scripts and Aakrosh Ratan provided early access to the YASRA assembler. Kandise Elliott worked her digital magic with the illustrations; Catherine Wardrop contributed the beautiful drawings of *Georgeantha* B. G. Briggs & L. A. S. Johnson and *Hypolaena* R. Br. Many thanks to Peter Stevens for having organized a highly stimulating symposium on plant diversification, to Peter Stevens and Victoria Hollowell for providing several helpful comments, and to Victoria Hollowell for her many contributions during editing.

² Department of Botany, University of Wisconsin, Madison, Wisconsin 53706, U.S.A. givnish@wisc.edu.

³ Department of Plant Biology, University of Georgia, Athens, Georgia 30602, U.S.A.

⁴ Department of Biological Sciences, University of Missouri, Columbia, Missouri 65211, U.S.A.

⁵ Department of Biology, Pennsylvania State University, University Park, Pennsylvania 16802, U.S.A.

⁶ Botanical Garden and Centre for Plant Research, University of British Columbia, Vancouver BC, Canada V6T 1Z4.

⁷ New York Botanical Garden, Bronx, New York 10458, U.S.A.

⁸ Botanic Gardens Trust, Sydney, New South Wales 2000, Australia.

⁹ Department of Biological Sciences, Northern Illinois University, DeKalb, Illinois 60115, U.S.A.

¹⁰ Department of Biology, Oberlin College, Oberlin, Ohio 44074, U.S.A.

¹¹ Department of Botany, University of Florida, Gainesville, Florida 32611, U.S.A.

¹² Florida Museum of Natural History, University of Florida, Gainesville, Florida 32611, U.S.A.

¹³ Western Australian Herbarium, Perth, Western Australia 6983, Australia.

doi: 10.3417/2010023

spread, and positive ecological feedback. Reproductive assurance in the absence of reliable animal visitation probably favored wind pollination in annuals and short-statured perennials of Centrolepidaceae in ephemerally wet depressions and windswept alpine sites.

Key words: Commelinids, correlated evolution, cyperids, graminids, long-branch attraction, molecular systematics, monocots, plastid, plastome, restiids, xyrids.

Monocots—with ca. 65,000 species in 82 families and 12 orders (Cameron et al., 2003; Givnish et al., 2006; Saarela et al., 2007; Angiosperm Phylogeny Group, 2009), and including such groups as the grasses, sedges, bromeliads, palms, ginger, bananas, orchids, irises, onions, asparagus, lilies, yams, pondweeds, aroids, and seagrasses—are one of the most diverse, morphologically varied, ecologically successful, and economically important clades of angiosperms. Since monocotyledons arose in the early Cretaceous (Herendeen & Crane, 1995; Bremer, 2000; Friis et al., 2004; Ramirez et al., 2007; Conran et al., 2009), they have radiated into almost every habitat on earth. Today, they dominate many terrestrial and aquatic ecosystems, display kaleidoscopic variation in vegetative and floral form, provide the basis for most of the human diet, support a huge horticultural industry, include large numbers of endangered taxa, and comprise nearly one fourth of all species and families of flowering plants. Understanding their origin, phylogeny, and patterns of morphological evolution, geographic diversification, and ecological radiation is thus a grand challenge and opportunity for evolutionary biologists.

Over the past 18 years, molecular systematics has revolutionized our understanding of monocot relationships (Chase et al., 1993, 1995a, b, 2000, 2006; Duvall et al., 1993a, b; Les et al., 1997; Givnish et al., 1999, 2005; Bremer, 2000, 2002; Kress et al., 2001; Hahn, 2002; Cameron et al., 2003; Michelangeli et al., 2003; Davis et al., 2004; Graham et al., 2006; Pires et al., 2006; Saarela et al., 2008). Such studies have led to a dramatic reclassification of the monocots (Angiosperm Phylogeny Group, 1998, 2003, 2009), with an ever-increasing understanding of relationships within and among the 12 orders currently recognized. Triumphs of monocot molecular systematics have included, first and foremost, the recognition of the commelinid clade (Chase et al., 1993, 1995a) composed of the orders Poales (the grasses, sedges, bromeliads, and their allies), Commelinales (the dayflowers, water hyacinths, and relatives), Zingiberales (the ginger, bananas, and related tropical monocots), Arecales (the palms), and the Australian Dasypogonales (Dasypogonaceae, Angiosperm Phylogeny Group, 2009, here recognized as an order following Givnish et al., 1999). This finding was buttressed by the demonstration that all five orders

share UV-fluorescent ferulic acid bound to cell walls (Harris & Hartley, 1980; Dahlgren et al., 1985; Rudall & Caddick, 1994; Harris & Tretheway, 2009) despite their otherwise great divergence in form and despite their relationships to each other that remain enigmatic based on molecular and morphological data (Chase, 2004; Graham et al., 2006). Other key advances have included the validation, to a large degree, of many of the orders inferred cladistically from morphology by Dahlgren et al. (1985); the identification of relationships among those orders; the placement of *Acorus* L. as sister to all other monocots (Duvall et al., 1993a); and the discovery that several genera, originally placed in Melanthiales based on morphology by Dahlgren et al. (1985) and Tamura (1998), actually belong to three other orders—including *Tofieldia* Huds. in Alismatales, *Narhecium* Huds. in Dioscoreales, and *Japonolirion* Nakai in Petrosaviales (Chase et al., 1995a, b, 2000, 2006; Zomlefer, 1999; Tamura et al., 2004). Perhaps most remarkably, Saarela et al. (2007) recently showed that highly reduced members of the aquatic family Hydatellaceae were not—as had long been believed (Hamann, 1976; Dahlgren et al., 1985; Angiosperm Phylogeny Group, 1998, 2003; Davis et al., 2004)—members of order Poales, and not even monocots, but were instead sister to waterlilies and other Nymphaeales, one of the earliest divergent clades of angiosperms (see Bosch et al., 2008, and Rudall et al., 2009, for corroborating morphological evidence).

Chase et al. (2006) provided the most powerful study of relationships across monocots to date, in terms of its combination of extensive taxon sampling (125 species stratified across 77 of 82 families) and sequence data per taxon (four plastid genes, two mitochondrial genes, one nuclear ribosomal gene). Based on their maximum parsimony (MP) analysis, *Acorus* is sister to all other monocots, followed by the successive divergence of Alismatales, Petrosaviales, Dioscoreales—Pandanales, Liliales, Asparagales, and the commelinids (Fig. 1). Based on Chase et al. (2006), all orders appear to be strongly supported, and all relationships among orders are resolved and—outside the commelinids—moderately to strongly supported. Yet, the MP strict consensus tree resulting from that study left many relationships within orders weakly supported, as well as a few of those among orders. Areas of substantial uncertainty include




Figure 1. Branching topology and support at the ordinal level in monocots, based on an analysis of four plastid genes, two mitochondrial genes, and one nuclear ribosomal gene (4777 informative sites, or 34.9 informative sites/taxon). Adapted from Chase et al. (2006).

relationships among many families within Poales, and among the five orders of commelinids; the positions of Asparagales and Liliales relative to the commelinids; and affinities of several families in Asparagales, Zingiberales, Dioscoreales, Pandanales, and Alismatales. In addition, relationships among commelinid orders found by Chase et al. (2006) weakly conflict with those identified by Graham et al. (2006) based on 16 kb of sequences for coding and noncoding plastid regions from 69 monocots representing 53 of 82 families. Research on relationships among various monocot groups has made enormous progress (see symposia volumes edited by Rudall et al., 1995; Wilson & Morrison, 2000; Columbus et al., 2006, 2007; Seberg et al., 2010). Yet much remains to be learned, perhaps because so many groups appear to have diverged more than 90 million years ago (Bremer, 2000, 2002; Givnish et al., 2000, 2005; Wikström et al., 2001; Janssen & Bremer, 2004), leaving only subtle traces of deep relationships in their current form and genetic sequences.

To address these uncertainties, develop a better understanding of broadscale relationships in monocots, and provide a strong basis for comparative studies of morphological, developmental, ecological,

and geographic differentiation in this important group, the National Science Foundation–funded Monocot AToL (Assembling the Tree of Life) project (<<http://www.botany.wisc./monatol/>>) is conducting a 5-year investigation of monocot evolution, building on the strong foundation of previous research by the international botanical community. Our aims are to: (1) develop a fully resolved, strongly supported, highly inclusive broadscale phylogeny for the monocots, based on sequencing transcriptomes of young leaf tissue for several dozen species, and whole-plastid genomes and targeted mitochondrial and nuclear genes for a few hundred species representing all monocot families and subfamilies; (2) score more than 200 morphological characters for the same extant species and 75 fossil taxa; (3) reassess the stratigraphy and classification of several fossil monocots and develop a new timeline for monocot evolution; (4) identify morphological synapomorphies to permit diagnosis of all major monocot clades, in both extant and fossil plants; (5) use the resulting phylogeny to conduct analyses of morphological, developmental, ecological, and biogeographic evolution using hundreds of taxa stratified across the monocots (e.g., see Davies et al., 2004; Givnish et al., 2005; Dunn et al., 2007); and ultimately (6) help train and inspire a new generation of monocot systematists in phylogenetics, genomics, and evolutionary biology. As a key part of this effort, we will sequence and analyze hundreds of complete plastid genome sequences. Almost all studies to date using DNA to infer relationships among monocots and other angiosperms have been phylogenetic, relying on sequences of only one or a few genes or gene spacers. Our approach will be phylogenomic, ultimately providing sequences for more than 100 chloroplast genes and the spacers between these genes, and generating an avalanche of new data with which to assess evolutionary relationships. This approach has proved fruitful in broadscale investigations of relationships among angiosperms and land plants (Leebens-Mack et al., 2005; Cai et al., 2006; Jansen et al., 2007; Moore et al., 2007; Wang et al., 2009). However, high data density per taxon must be complemented with a reasonably dense stratification of taxon sampling across the range of organisms being investigated in order to minimize errors in phylogenetic inference due to long-branch attraction (e.g., see Goremykin et al., 2003, vs. Soltis & Soltis, 2004; Leebens-Mack et al., 2005).

In sequencing plastomes and targeted mitochondrial and nuclear genes, we are collaborating closely with members of the Angiosperm AToL team (<<http://www.flmnh.ufl.edu/angiospermATOL/>>), who have already sequenced more than 30 plastomes and several mitochondrial and nuclear genes for nonmonocots.

We are also coordinating our efforts with members of the European monocot initiative (supported by a grant from the Leverhulme Trust) who are sequencing two plastid genes (*rbcL* and *matK*) for representatives of all ca. 2400 monocot genera. These efforts are complementary, in taxonomic coverage for the Angiosperm AToL, and in our resolution of deep monocot nodes while the Europeans link our backbone phylogeny to the remaining monocot genera. We are also collaborating with the 1000 Plants Initiative (<http://www.onekp.com>) to sequence shoot transcriptomes for several dozen species. We believe that the unparalleled amount of genetic information from all three plant genomes obtained in our AToL project should provide the most powerful analysis of relationships for any major plant group studied to date.

In this paper, we present a demonstration of the utility of a plastome-based phylogenomic approach, focusing on relationships among families of Poales and their immediate relatives among the commelinids. Poales is the second largest order of monocots, and economically surely the most important, as a consequence of its including the cereal and pasture grasses, which contribute so much to the human diet, directly or indirectly; the bamboos, a key source of building materials; and the restioids, sedges, rushes, cattails, and bur reeds, which form such key components of wetland ecosystems locally and worldwide. Poales and the commelinids include many of the nodes in the monocot tree of life that have proven most difficult to resolve using traditional phylogenetic techniques. Here we present a new, plastome-based phylogeny for the monocots and discuss its implications for relationships among families of Poales and its immediate commelinid relatives. We then use this phylogeny to analyze broadscale evolutionary patterns within Poales, including habitat diversification and, especially, the pattern and potential causes of multiple origins of wind pollination, an otherwise fairly uncommon mechanism in the monocots. We conclude with a brief discussion of some of the limitations of plastome-based phylogenomics for understanding monocot phylogeny and evolution.

METHODS

TAXON AND GENE SAMPLING

The 83 taxa included in our study represent most major clades of angiosperms sensu Angiosperm Phylogeny Group (2009) (Appendix 1). New draft plastid genome sequences were generated for 44 of the 83 taxa included in the study (Appendix 1). These new data greatly increase the number of monocot sequences included in plastid phylogenomic analyses.

Sixty-four species acted as placeholders for 11 monocot orders and 32 of 82 monocot families. Taxon sampling is concentrated on Poales and, to a lesser extent, its putative relatives among the commelinids and Asparagales. Commelinids sequenced included members of 15 of 16 families of order Poales (Anarthriaceae not sampled); one of five families of Commelinales; two of eight families of Zingiberales; and the single families in Arecales and Dasygogonales (Dasygogonaceae sensu Angiosperm Phylogeny Group, 2009). Nonmonocot angiosperms included representatives of an additional 18 orders of flowering plants (Appendix 1). We used *Amborella* Baill., the consensus sister taxon to all other angiosperms (Jansen et al., 2007; Moore et al., 2007) as the outgroup.

PLASTOME SEQUENCING

We used next-generation sequencing to generate 44 plastid genome sequences (Appendix 1) using one of two strategies. Following methods described by Jansen et al. (2005), plastids of *Lilium* L., *Hosta* Tratt., *Tradescantia* L., *Brocchinia* Schult. f., *Neoregelia* L. B. Sm., *Pitcairnia* L'Hér., and *Puya* Molina were isolated on a sucrose gradient and used as templates for rolling circle DNA amplification. Plastome-enriched amplicons were sequenced on a Roche GS-FLX sequencer (Roche, Branford, Connecticut, U.S.A.) using 454 pyrosequencing technology (Moore et al., 2007). The remaining 37 draft plastome sequences were assembled from whole-genome shotgun sequences generated on an Illumina GS sequencer (Illumina, Inc., San Diego, California, U.S.A.). The proportion of plastid DNA in isolated DNAs was estimated through quantitative real-time PCR amplification of a 150-bp portion of the *rbcL* gene. Samples estimated to have at least 5% plastid DNA were prepared for whole-genome shotgun sequencing on the Illumina sequencing platform, following the manufacturer's protocol. Bar codes were ligated to templates, and at least one million 75-bp reads were generated for each taxon. De novo assemblies of sequences generated on the 454 platform were constructed using the manufacturer's Newbler assembler (<http://www.454.com>) and the MIRA assembler (http://www.chevreux.org/projects_mira.html; Chevreux et al., 2004). Resulting assemblies were inspected using Consed (Gordon et al., 1998) and Sequencher (<http://www.genecodes.com/>). Genes were annotated and gene sequences were extracted from assemblies using the DOGMA webserver (<http://dogma.cbb.utexas.edu>), Wyman et al., 2004, and see below). Sequences generated using the Illumina short-read platform were subjected to reference-based assembly using the YASRA assembler (Ratan, 2009;

download available with documentation at http://www.bx.psu.edu/miller_lab) to layer short reads on a reference genome while allowing substantial sequence divergence (< 85% sequence identity). For each taxon, the most closely related plastome genome sequence available in GenBank was initially used as the reference genome. In addition, our draft genome sequences were used as alternative references for assembly of related genomes. For example, the *Cyperus alternifolius* L. draft plastome was used as a reference for *Juncus effusus* L. and *Thurnia sphaerocephala* Hook. f. plastome assembly and vice versa. Final assemblies for each taxon were compiled in Sequencher. As described above, genes were annotated and extracted from the resulting contigs using DOGMA. DOGMA identifies genes through BLASTX searches against a database of amino acid sequences extracted from exemplar plastid genomes. As a consequence, gaps in the assemblies, rare sequencing errors, and assembly errors that introduce frame shifts or stop codons may result in annotation of gene fragments rather than full-length genes. Only full- or near-full-length gene annotations were included in alignments and phylogenetic analyses. Therefore, individual taxa may be missing genes due to low sequencing coverage, assembly errors, or evolutionary loss (e.g., the loss of the *yef2* gene in the inverted repeat [IR] region of the plastome on the branch leading to Poaceae/Scitaceae/Joinvilleaceae). The only taxa with substantial numbers of missing genes were those with less than 15 \times coverage under next-generation sequencing, including *Abolboda macrostachya* Spruce ex Malme, *Mayaca fluviatilis* Aubl., and *Syngonanthus chrysanthus* Ruhland, and *Joinvillea plicata* (Hook. f.) Newell & B. C. Stone under ordinary direct sequencing. All gene sequences included in alignments and phylogenetic analyses have been deposited into GenBank. All analyses were based on the exons of 77 protein-coding plastid genes and four ribosomal RNAs (rRNAs), with a total of 109,134 aligned bases. This set of 81 plastid genes was also used by Jansen et al. (2007) to resolve broadscale relationships across flowering plants.

SEQUENCE ALIGNMENT

Following Jansen et al. (2007), 81 plastome-encoded gene sequences were aligned individually and alignments concatenated into a single nexus file for phylogenetic analyses; the file is posted at the Monocot AToL project website (see <http://chloroplast.cbio.psu.edu/supplement.html>). Perl scripts were written to sort gene sequences extracted from DOGMA annotations for each taxon into multitaxon fasta files for each gene, to align gene sequences using MUSCLE

(Edgar, 2004), and to concatenate alignments in a Nexus file. Missing genes were filled with Ns in the concatenated alignment.

PHYLOGENETIC ANALYSES

We inferred relationships among taxa from the nucleotide data using MP and maximum likelihood (ML). MP analyses were run using PAUP* 4.0d102 (Swofford, 2002). Individual nucleotides were considered to be multistate, unordered characters of equal weight; unknown bases were treated as uncertainties. Gapped cells were treated as missing data and we did not attempt to score indels. To evaluate the possibility of multiple islands of equally most parsimonious trees (Maddison, 1991), we ran heuristic searches seeded with 100 random-addition sequences, employing tree bisection-reconnection (TBR) swapping while retaining up to 100 trees per iteration. Bootstrap analysis (Felsenstein, 1985) was used to assess the relative support for each node in the single shortest tree found, using 200 random resamplings of the data and retaining up to 100 trees per resampling. Consistency indices, including autapomorphies (CI) and excluding them (CI'), were calculated to measure the relative extent of homoplasy in the data (Givnish & Sytsma, 1997).

ML analyses were performed on concatenated alignments using RAxML version 7.0.4 (Stamatakis, 2004). A single substitution process was modeled as general time reversible (GTR) plus gamma for the entire concatenated alignment (i.e., data were not partitioned). Among-site variation in substitution rates was modeled using the discrete approximation of the gamma distribution (Yang, 1994) with 25 rate classes. The ML bootstrap analysis included 250 pseudo-replicates drawn from the concatenated alignment.

ANCESTRAL HABITAT RECONSTRUCTION

To trace patterns of habitat evolution within the order Poales, we overlaid various environmental characteristics on the commelinid portion of the ML cladogram, simplified to the family level, using parsimony as implemented in MacClade 4.0 (Maddison & Maddison, 1992) to infer ancestral states, resolving all of the most parsimonious states at each node. Given their distinctive ecologies, we separated subfamily Anomochlooideae from the other Poaceae sampled (a subset of the bistigmatic clade = subfamily Puelioideae + the PACCMAD-BEP clade of grasses; the latter consists of Panicoideae, Arundinoideae, Chloridoideae, Centothecoideae, Micrairoideae, Aristidoideae, Danthonioideae-Bambusoideae, Ehrhartoideae, and Pooideae (Sánchez-Ken et al., 2007). We interpolated the subfamily Phar-

oideae (unsampled in our phylogenetic study) between Anomochlooideae and Puelioideae, given that recent molecular studies place them there with high support, and given the distinctive ecology of pharoids and puelioids versus most members of the huge PACC-MAD-BEP clade, which includes 99% of all grass genera and species (Soderstrom & Calderón, 1971; Clark et al., 1995, 2000; Grass Phylogeny Working Group I, 2001; Hodkinson et al., 2007a, b; Boucheanak-Khelladi et al., 2008). Characteristics overlaid include: (a) light availability (sunny vs. shady); (b) moisture supply (soil wet or inundated vs. well drained or dry); (c) soil fertility (highly infertile [e.g., sand or sandstone] vs. fertile); and (d) fire prevalence (high vs. low/absent). Data on ecology were drawn from summaries in Dahlgren et al. (1985), Givnish et al. (1999, 2000, 2004, 2007), Linder and Rudall (2005), and Sokoloff et al. (2009). We used the ML phylogeny, including Poales and families of the other commelinids studied as ingroups and *Apostasia* Blume of Orchidaceae as the outgroup, because it resolves a few crucial nodes differently and with far higher support than the MP tree, and because ML generally is less susceptible to problems caused by extensive change in evolutionary rates over time (i.e., heterotachy) and long-branch attraction (e.g., Huelsenbeck & Hillis, 1993; Huelsenbeck, 1995; Chang, 1996; Swofford et al., 2001; Gadagkar & Kumar, 2005; Leebens-Mack et al., 2005; Jansen et al., 2007; Moore et al., 2007; Whitfield & Lockhart, 2007; Wang et al., 2009). Where necessary in large, ecologically diverse groups represented by single taxa, we overlaid the ancestral conditions previously inferred for such families based on detailed infrafamilial molecular studies (e.g., Givnish et al., 2000, 2004, 2007) or inferred what those ancestral conditions would have been, given the habitats of the first several earliest-divergent lineages within those families (Appendix 2).

ORIGINS OF WIND POLLINATION AND PATTERNS OF CORRELATED EVOLUTION

We also overlaid wind versus animal pollination on the simplified Poales tree using MacClade, following earlier analyses by Givnish et al. (1999) and Linder and Rudall (2005). We drew data from those papers and from Newell (1969), Soderstrom and Calderón (1971, 1979), Henderson (1986), Stützel (1986, 1990), Listabarth (1992), Soreng and Davis (1998), Rosa and Scatena (2003, 2007), Blüthgen et al. (2004a, b), Blüthgen and Fiedler (2004), Ramos et al. (2005), Moura et al. (2008), Oriani et al. (2009), and Sokoloff et al. (2009) (see Appendix 2, Table A2). Previous authors have argued that wind pollination should be

positively associated with visually inconspicuous, unisexual flowers; an absence of nectar; single ovules; growth in open or seasonally open, windy environments; and a taxon's local abundance (see Faegri & van der Pijl, 1979; Regal, 1982; Cox, 1991; Linder, 1998; Weller et al., 1998; Givnish et al., 1999; Cully et al., 2002; Friedman & Barrett, 2008). To evaluate how these ideas apply to Poales, we conducted formal tests of correlated evolution of pollination mechanism (wind vs. animal) with plant habitat, nectaries (presence or absence), floral sexuality (hermaphroditic vs. unisexual flowers), sexual system (cosexuality vs. dioecy), ovule number (one vs. many), floral size (small vs. large petals or other visual displays), and floral showiness (nonshowy vs. showy colors) using the BayesDiscrete module of BayesTraits (available at <<http://www.evolution.rdg.ac.uk>>). BayesDiscrete tests for significant patterns of correlated evolution of two binary traits by comparing the fit (ln likelihood) to models that assume their independent versus dependent evolution on the phylogeny provided using continuous-time Markov models (Pagel & Meade, 2006). For each trait, we conducted two sets of nested ML analyses, either setting the branch-length parameter κ equal to 1 or allowing it to assume its optimal value under ML analysis. Values of $\kappa < 1$ result in a scaling that reduces the length of longer branches more than shorter branches. Our approach directly parallels that used by Friedman and Barrett (2008) to test for correlated evolutionary patterns involving wind pollination across angiosperms. We used the nonsimplified commelinid portion of our ML tree for topology and branch lengths. Character states for all taxa in the ML tree (see Appendix 2) were drawn from the pollination references previously cited and from two data sets kindly provided by Jannice Friedman, Jana Vamosi, and Spencer Barrett, and by George Weiblen. We scored the highly unusual (if not, indeed, unique) case of *Chamaedorea* Willd. in which thrips are involved in releasing clouds of pollen that move from staminate plants to pistillate plants via the wind (Listabarth, 1992), as animal pollinated, given the role that animals play in this case of animal-induced wind pollination.

RESULTS

PHYLOGENY

MP yielded a single, fully resolved tree of length 152,366 steps (Fig. 2). Overall, there were 25,107 parsimony-informative characters, of which 22,156 were informative within the monocots. Across all taxa, 12,634 characters were variable but uninformative and 71,834 were constant. Monocots were monophyletic and




Figure 2. Phylogram for the single MP tree resulting from analysis of the plastome data, rooted using *Amborella* Baill. Branch lengths are proportional to the number of inferred substitutions down each branch. Bootstrap support for each node is shown above the corresponding branch. Monocots are highlighted with magenta branches; cyperids, xyrids, restiids, and graminids, with colored boxes. Full scientific names and authorities for the exemplars shown are given in Appendix 1.

Acorus was sister to all other taxa, both with 100% bootstrap support. At the ordinal level, Alismatales (*Lemna* L.), Liliales (*Lilium*), Dioscoreales–Pandanales (*Dioscorea* L.–*Pandanus* Parkinson), and finally Asparagales and the commelinids diverged from progressively more recent nodes. Bootstrap support for the branching sequence of Liliales, Dioscoreales–Pandanales, and Asparagales was, however, weak (52%–56%), as was that for the monophyly of Asparagales (Fig. 2). The

commelinids as a whole, and each of their five orders, had 100% bootstrap support. Poales was resolved as sister to Arecales, and Dasygogonales as sister to Commelinales–Zingiberales, but both relationships had weak support.

Within Poales, Bromeliaceae were sister to all other taxa, with 100% bootstrap support, followed by the divergence of Typhaceae and Rapateaceae at the two succeeding nodes, with 97% and 100% support,

respectively (Fig. 2). The MP tree recovered the xyrids (Eriocaulaceae, Mayacaceae, Xyridaceae) as monophyletic with 68% bootstrap support and placed them sister to the cyperids (Cyperaceae, Juncaceae, Thurniaceae) with 78% bootstrap support. Within the xyrids, *Mayaca* Aubl. was placed sister to *Syngonanthus* Ruhland of Eriocaulaceae with 99% bootstrap support. Branches among the cyperids all had 100% bootstrap support. The restiids (Restionaceae, Centrolepidaceae, and Anathriaceae, with the last small family unsampled here) were monophyletic, with 100% bootstrap support, and were sister (with 94% bootstrap support) to the graminid clade of grasses and their allies. Within the graminids, *Flagellaria* L. was sister to all other taxa with 100% support; *Joinvillea* Gaudich. ex Brongn. & Gris was sister to the remaining taxa, also with 100% support. Finally, Ecdeiocolaceae was resolved as monophyletic (100% support) and sister to the grasses (94% support).

Across monocots, inferred branch lengths were especially short in Acoraceae, Arecaceae, Bromeliaceae, and Typhaceae, and especially long in the graminids, restiids, xyrids, and cyperids (Fig. 2). Seven of these eight clades are part of the commelinids. Relative to the commelinid crown-group root node, branch lengths averaged 1734 ± 170 (mean \pm SD) steps for palms, 2112 ± 103 for bromeliads, and 2701 ± 67 for cattails and bur-reeds, compared with 7027 ± 1062 for xyrids, 7821 ± 1102 for cyperids, 8644 ± 1210 for restiids, and 9152 ± 1416 for graminids. Rates of plastid sequence evolution thus vary by as much as 5.2-fold within the commelinids, based on MP analysis.

ML produced a single, fully resolved, well-supported phylogeny (Fig. 3). This tree is strikingly similar in most regards to the MP tree, with four key exceptions. First, *Lilium* (Liliales) was resolved as sister to the Asparagales–commelinid clade; this position has 94% bootstrap support. Second, Asparagales had 100% bootstrap support as being monophyletic and 99% support as being sister to the commelinids in the ML tree, versus 56% for both conditions in the MP tree. Support for the topology of the ML tree was generally higher than that for the MP tree, with 63 of 67 nodes within the monocots having $\geq 94\%$ bootstrap support, and all 18 nodes outside the monocots having 100% support (Fig. 3). Third, ML provided a different and much more strongly supported resolution of the commelinid orders, with Poales being sister to Commelinales–Zingiberales with 93% bootstrap support, and Arecaceae being sister to Dasypogonales with 86% bootstrap support. Finally, in the ML tree, the xyrids formed a grade, not a clade, and were associated with the graminid–restiid clade, not the cyperids. *Abolboda* Bonpl. of Xyrida-

ceae was sister to the restiid–graminid clade with 100% bootstrap support; *Mayaca* of Mayacaceae and *Syngonanthus* of Eriocaulaceae were sister to *Abolboda* plus the restiid–graminid clade (Fig. 3).

ANCESTRAL HABITAT RECONSTRUCTION

We infer that the ancestral Poales occupied habitats that were sunny, wet, possibly nutrient poor, and fireswept (Figs. 4, 5), much like the conditions occupied by *Brocchinia prismatica* L. B. Sm., *B. melanacra* L. B. Sm., *B. reducta* Baker, *Lindmania guianensis* (Beer) Mez, and other early divergent members of Bromeliaceae today (Givnish et al., 1997, 2007). Low-nutrient conditions characterize the ancestral Poales only under ACCTAN. However, infertile soils are more likely than fertile soils to favor fire (Givnish, 1980), the inferred condition for ancestral Poalean habitats (Fig. 5B). Both ACCTAN and DELTAN reconstruct infertile soils as the ancestral state for the clade including Rapateaceae and its sister (Fig. 5A). Open habitats typify most of the early divergent families—from bromeliads to the restiids—although bromeliads and rapateads include a number of reinvasions of shaded sites (Givnish et al., 2000, 2004, 2007), as do sedges (e.g., in *Becquerelia* Brongn. and *Carex* L.) and rushes (e.g., in *Luzula* DC.), which can be inferred from the phylogenies provided by Drábková and Vlček (2009) and Muasya et al. (2009). Shady habitats were invaded by members of a grade running from *Flagellaria* and *Joinvillea* through the anomochlooid and pharoid grasses, with reinvasions of sunny habitats in Ecdeiocolaceae and the PACCMAD-BEP grasses (Fig. 4A). Damp or wet soils appear to be the ancestral condition in Poales from bromeliads (e.g., *Brocchinia*, *Lindmania* Mez) through at least Xyridaceae (*Abolboda*), with a transition to well-drained rainforest soils for *Flagellaria*, *Joinvillea*, Anomochloideae, and Pharioideae (Fig. 4B). Several Centrolepidaceae and Restionaceae grow in permanently or seasonally inundated soils, but others occupy well-drained sites.

If we assume that the ancestral habitat of Poales had highly infertile soils (see above), then such soils would have characterized all families and ancestors from Bromeliaceae through the restiids, with the exception of Typhaceae, Cyperaceae, and Juncaceae on richer soils, although some members of the latter two families now occupy highly infertile substrates as well. The terminal clade of Poales (Flagellariaceae through Poaceae) typically occupies more fertile substrates, with the exception of Ecdeiocolaceae. Finally, fireswept habitats characterized families from Bromeliaceae through Xyridaceae and possibly the restiids, with such disturbance being lost in the aquatic *Thurnia* Hook. f. (but perhaps not in *Prionium* E. Mey.), the aquatic *Mayaca*, the aquatic and alpine




Figure 3. Phylogram for the single ML tree. The nonmonocot portion of the tree is not shown. The resolution of that portion is, however, identical to that shown in Figure 1, with 100% bootstrap for all nodes. Bootstrap support for each node within the monocots is shown above the corresponding branch. Cyperids, xyrids, restiids, and graminids are highlighted with colored boxes.




Figure 4. Inferred evolution of (A) light availability and (B) moisture supply in Poales and immediate outgroups at the family level under parsimony. Cases in which the ancestral character state for a family could not be inferred unequivocally (see text) were coded as polymorphic, and both character states for the current-day, terminal taxa are indicated in the split box. Gray branches indicate equivocal resolution of character states in ancestral taxa.

Centrolepidaceae, and the forest-inhabiting grade from *Flagellaria* through the early divergent grass families, with fire ecology recurring in Ecdeiocoleaceae and Poaceae (Fig. 5B). If Anarthriaceae had been included in this analysis, sister to the other restiid subfamilies as in other recent analyses (e.g., Chase et al., 2006), then Restionaceae would have been resolved as retaining fireswept habitats rather than invading them anew.

MULTIPLE ORIGINS OF WIND POLLINATION AND CORRELATED TRAITS

Ancestral character-state reconstruction using the ML tree implies that wind pollination evolved at least

five times in Poales: in Typhaceae, cyperids, restiids, Ecdeiocoleaceae, and the PACCMAD-BEP clade of Poaceae (Fig. 6). Animal pollination is inferred to be homologous across the commelinids.

Based on ML testing in BayesTraits, pollination mechanisms in Poales and its immediate relatives showed correlated evolution with (1) floral showiness, (2) flower size, (3) floral sexuality, (4) nectar production, and (5) habitat openness (Table 1). Wind pollination was, as expected, associated with smaller, less conspicuous, often unisexual flowers and with open habitats. In Poales and its immediate relatives, there was no evidence for correlated evolution between pollination mechanism and ovule number or sexual system (cosexuality vs. dioecy); the latter finding




Figure 5. Inferred evolution of (A) nutrient supply and (B) fire frequency in Poales and immediate outgroups at the family level (see Fig. 4).

implies that unisexual flowers associated with wind pollination are typically found in cosexual plants.

DISCUSSION

PHYLOGENY

ML analysis of 81 coding regions from the plastid genome produced—within the limits of taxon sampling—the best resolved and most strongly supported phylogeny to date of Poales, the commelinids, and the monocots as whole, with a mean bootstrap support of 98.2% for all monocot nodes (Fig. 3). We focus on the ML tree because it had much higher bootstrap support for several crucial nodes within and outside Poales than the MP tree, including some that differ between the two trees; because Poales exhibits striking rate heterogeneity and a combination of very short and

very long branches in close proximity; and because ML generally outperforms MP in reconstructing phylogenies when faced with rate heterogeneity and long-branch attraction. The nodes at which the ML and MP trees differ are just those with very short and very long branches immediately juxtaposed, where long-branch attraction might be expected to be especially likely to distort phylogenetic reconstruction via parsimony. This was seen in the placement of the xyrids within Poales, the monophyly of the xyrids, the sister group to the Poales, relationships among the other commelinid orders, and the placement of Liliales (Fig. 3). The much higher rates of evolutionary divergence of plastid coding regions in Poaceae and allied families versus Arecaceae and Bromeliaceae have long been recognized (Gaut et al., 1992, 1996; Givnish et al., 1999, 2004; Smith & Donoghue, 2008).




Figure 6. Minimum of five independent origins of wind pollination within Poales and immediate outgroups, plotted at the family level. Drawings show typical floral and/or inflorescence form for each origin. From top to bottom: *Tripsacum dactyloides* L. (Poaceae), staminate and pistillate portions of inflorescence; *Georgeantha hexandra* B. G. Briggs & L. A. S. Johnson (Ecdeiocoleaceae), floral spike with zones of male and female flowers; *Hypolaena fastigiata* R. Br. (Restionaceae), staminate and pistillate inflorescences borne on separate plants; *Carex pilulifera* Willd. ex Kunth (Cyperaceae), staminate and pistillate portions of inflorescence; and *Sparganium emersum* Rehm. (Typhaceae), staminate and pistillate portions of inflorescence. Note long and/or plumose branches of styles in each case and mostly large, dangling (often centrally attached) anthers.

Table 1. Summary of tests for correlated evolution of wind pollination with various plant traits.

Trait	Significance with $\kappa = 1$	Significance with optimal κ
Flower size	$P < 0.012$	$P < 0.027$
Floral showiness	$P < 0.0034$	$P < 0.013$
Floral sexuality	$P < 0.0002$	$P < 0.0005$
Habitat openness	$P < 0.013$	$P < 0.02$
Nectar production	$P < 0.035$	$P < 0.038$
Dioecy	NS	NS
Ovule number	NS	NS

NS, not significant.

RELATIONSHIPS WITHIN POALES

The ML and MP trees agree in placing, with strong support, Bromeliaceae, Typhaceae, and Rapateaceae as successive sister lineages to all other members of order Poales (Figs. 2, 3). The lowest support, with 87%–97% bootstrap values, is for the position of Typhaceae, whose sister group has the shortest branch for any family within Poales. Resolving the relationships among these three families has been problematic. Michelangeli et al. (2003) used sequences of plastid *rbcL* and mitochondrial *atpB* to place Rapateaceae plus the xyrids as sister to all other Poales. Bremer (2002) and Davis et al. (2004), using the same genes, placed Rapateaceae alone as sister to all other Poales, with Bromeliaceae being sister to Typhaceae, and both being sister to the remainder of the order, followed by Eriocaulaceae plus most Xyridaceae; jackknife support for most of these relationships, however, was quite low. Givnish et al. (2005, 2006) used *ndhF* sequences to place Bromeliaceae plus Typhaceae sister to all other Poales, followed by Rapateaceae. Christin et al. (2008) used *ndhF* and *rbcL* to reach a similar conclusion, but did not include Rapateaceae. Graham et al. (2006) used sequences of 17 plastid genes and spacers to place Typhaceae as sister to all other Poales, followed by Bromeliaceae. Only Chase et al. (2006), using data for four plastid genes, two nuclear genes, and one mitochondrial gene, recovered the branching pattern for Bromeliaceae, Typhaceae, and Rapateaceae that we found, but they had low (less than 50%) bootstrap support for the position of Typhaceae. The strong support for the branching pattern of all three families relative to other Poales in both our analyses should settle their position and branching pattern. The absence of septal nectaries from all Poales except Bromeliaceae and a few derived Rapateaceae (where such nectaries have apparently reevolved [Givnish et al., 2000]) helps support the placement of Bromeliaceae at the base of the order. ML placed the cyperids sister to all other

Poales except Bromeliaceae, Typhaceae, and Rapateaceae, with the branching topology among the three families identical to that found by most broadscale monocot analyses since Givnish et al. (1999) used *rbcL* sequences to place Thurniaceae sister to Cyperaceae and Juncaceae.

The most vexing issue involves the position of the xyrids. Our MP tree identified these as monophyletic and sister to the cyperids, but it identified *Abolboda* of Xyridaceae as sister to *Mayaca* of Mayaceae and *Syngonanthus* of Eriocaulaceae (Fig. 2). However, there was only 68% bootstrap support for the monophyly of the xyrids in the MP tree, and only 78% support for its position sister to the cyperids. In the ML tree (Fig. 3), the xyrids were a grade rather than a clade, with *Abolboda* sister to the restiids–graminids with 100% support, followed by *Mayaca*–*Syngonanthus* with 97% support. The difficulty here, undoubtedly, is the great length of the branches leading to each of the three families of xyrids, relative to the short length of branches joining those families. Our placement of the xyrid families, based on the ML tree, should be viewed as tentative, with a need for sequencing additional species to help break up the long branches within each family. Givnish et al. (2005, 2006) and Chase et al. (2006) both placed Eriocaulaceae and Xyridaceae sister to each other in analyses that included *Mayaca*; Bremer (2002) placed these families sister to each other but did not include *Mayaca*. Givnish et al. (2005, 2006) placed Eriocaulaceae–Xyridaceae sister to the cyperids, with Mayaceae sister to the broader group; Chase et al. (2006) identified *Mayaca* as sister to the cyperids, followed by Eriocaulaceae–Xyridaceae. Graham et al. (2006) obtained results consistent with those of Givnish et al. (2005, 2006), but did not include Eriocaulaceae. Davis et al. (2004) placed *Mayaca* plus *Xyris* L. (plus *Hydatella* Diels) sister to the cyperids, with the remainder of Xyridaceae plus Eriocaulaceae sister to that group. Givnish et al. (1999), Bremer (2002), and Christin et al. (2008) placed Eriocaulaceae–Xyridaceae sister to the restiid–graminid clade. We note that that position is consistent with Eriocaulaceae and Xyridaceae sharing the absence of a parietal cell in the nucellus with the graminids, but excluding *Flagellaria* (Rudall, 1997) and restiids (Givnish et al., 1999). Missing data for the xyrids (see above) do not account for their placement. When we exclude all genes not represented in all three xyrid placeholders, ML yields exactly the same branching topology (albeit with weaker support) within Poales. Placement of the xyrids should not be viewed simply as a choice between being sister to the cyperids versus the restiid–graminid clade. When we included *Mayaca* while excluding *Abolboda* and *Syngonanthus* in our

own analyses, *Mayaca* was sister to the cyperids plus the restiid–graminid clade. Christin et al. (2008) placed *Mayaca* sister to the cyperids alone.

Morphologically, there is relatively little evidence tying Mayacaceae to Eriocaulaceae and Xyridaceae (see Givnish et al., 1999; Stevens, 2009). These three families are the remnants of the order Commelinales recognized by Dahlgren and Clifford (1982) and Dahlgren et al. (1985) before Givnish et al. (1999) used *rbcL* sequence data to exclude Commelinaceae and Rapateaceae. All five families share nuclear endosperm and showy flowers with differentiated petals and sepals; both characters are more broadly plesiomorphic in commelinids (Givnish et al., 1999; Stevens, 2009). In other words, there is relatively little evidence to support an expectation that these three families will form a clade when additional plastome sequence data are added to future analyses.

The position of the restiids as sister to the graminids in both the MP and ML trees (Figs. 2, 3) accords with the results presented by Chase et al. (2006) and is consistent with their positions in the incompletely resolved tree presented by Givnish et al. (2005, 2006). The only point of disagreement involves the placement of *Flagellaria*, which was sister to the restiids and the graminids by Michelangeli et al. (2003), sister to the cyperids plus *Xyris*, *Mayaca*, and *Trithuria* Hook. f. (Hydatellaceae) by Davis et al. (2004), and sister to *Elegia* L. (Restionaceae) plus representatives of the other graminid families by Givnish et al. (1999) and Graham et al. (2006). Our sampling includes only a single representative for each of the families Restionaceae and Centrolepidaceae, with none for Anarthriaceae. Anarthriaceae, when included in broadscale phylogenetic studies based on molecular data, has usually been placed sister to Centrolepidaceae–Restionaceae (Briggs et al., 2000; Bremer, 2002; Michelangeli et al., 2003; Davis et al., 2004; Chase et al., 2006). Centrolepidaceae was sister to Restionaceae in most broadscale molecular studies, but some (e.g., Bremer, 2002; Briggs et al., 2010) have embedded it in Restionaceae, while Briggs and Linder (2009) regarded its position as unresolved.

Within the graminids, our data confirm that Flagellariaceae, Joinvilleaceae, and Ecdeiocoleaceae are successively sister to ever-narrower subsets of the graminids, with Ecdeiocoleaceae sister to Poaceae. This branching topology concurs with those documented by Bremer (2002), Chase et al. (2006), and Graham et al. (2006), but conflicts with those of Marchant and Briggs (2007), Christin et al. (2008), and others that place *Joinvillea* sister to Ecdeiocoleaceae, with both sister to Poaceae. Early studies based on morphological or molecular data either

assumed or concluded that *Joinvillea* was sister to the grasses (e.g., Campbell & Kellogg, 1987; Chase et al., 1995a, b; Clark et al., 1995; Kellogg & Linder, 1995; Stevenson & Loconte, 1995; Soreng & Davis, 1998; Grass Phylogeny Working Group I, 2001). Briggs et al. (2010) were unable to resolve a trichotomy involving the grasses, *Joinvillea*, and Ecdeiocoleaceae. Doyle et al. (1992) discovered a 28-kb inversion in the chloroplast genome that united Joinvilleaceae, Ecdeiocoleaceae, and Poaceae; they detected a 6-kb inversion in Joinvilleaceae and Poaceae, but were unable to amplify the region in question for Ecdeiocoleaceae. Several subsequent authors confused the absence of data on the 6-kb inversion with absence of that inversion in Ecdeiocoleaceae, until Michelangeli et al. (2003) demonstrated that the inversion was indeed present in that family, and that *rbcL* and *atpB* sequence data supported (albeit quite weakly) a sister-group relationship between *Ecdeiocolea* F. Muell. and the grasses. In retrospect, a key morphological character used by Campbell and Kellogg (1987), Kellogg and Linder (1995), and Kellogg (2000) as a synapomorphy to unite Joinvilleaceae and Poaceae—namely, the presence of long and short cells in the leaf epidermis—may be seen to exclude Ecdeiocoleaceae. Although adult plants of *Ecdeiocolea* have scarcely any development of leaf blades, juvenile plants of *Ecdeiocolea* and adult *Georgeantha* B. G. Briggs & L. A. S. Johnson (Briggs & Johnson, 1998) show substantial leaf blades with no evidence of a long and short cell pattern (Briggs, pers. obs.). Thus, the presence of long and short cells in the leaf epidermis either evolved twice (in Poaceae and Joinvilleaceae) or—perhaps more likely—arose once in the common ancestor of Joinvilleaceae–(Ecdeiocoleaceae–Poaceae) and subsequently was lost in the highly reduced Ecdeiocoleaceae.

Within Poaceae, our data confirm that *Anomochloa* Brongn. and *Streptochaeta* Schrad. ex Nees are sister taxa, forming subfamily Anomochlooideae; this subfamily is, in turn, sister to the remainder of the family (see Clark et al., 1995; Grass Phylogeny Working Group I, 2001; Christin et al., 2008). The latter lineage is characterized by the presence of the typical grass-type spikelet found in all grasses except Anomochlooideae, and is therefore termed the spikelet clade (Grass Phylogeny Working Group I, 2001). Relationships among the remaining grasses generally accord with those found using molecular data by Grass Phylogeny Working Group I (2001), Hodkinson et al. (2007a, b), Bouchenak-Khelladi et al. (2008), and Saarela and Graham (2010). We hope to add several additional plastome sequences in the near future to permit detailed analyses of evolution within the Poaceae.

RELATIONSHIPS AMONG COMMELINID ORDERS

The ML phylogeny presented provides the first well-supported (93% bootstrap) evidence for the sister group to Poales, identifying the Commelinales–Zingiberales clade as the closest relative to the grasses, bromeliads, and their allies (Fig. 3). The two remaining commelinid orders, Arecales and Dasypogonales, also form a clade with relatively strong (87% bootstrap) support, which is sister to the clade formed by the previous three orders. None of the recent analyses of broadscale relationships among monocots found this branching topology. Givnish et al. (1999) and Davis et al. (2004) came closest, placing Arecales or Dasypogonales sister to Commelinales–Zingiberales plus Poales. Given that our MP analysis retrieved an alternative, albeit weakly supported topology (Fig. 2), our conclusions regarding ordinal relationships among commelinids should be viewed somewhat tentatively. We plan to include multiple representatives of each family in analyses in the near future, using increased taxon sampling density to confront the difficulties caused by long-branch attraction and striking variation in evolutionary rates within the commelinids. For now, however, we note that the sister-group relationship between Poales and Commelinales–Zingiberales is supported by possession of (1) starchy endosperm across all taxa surveyed and (2) a distichous or tristichous phyllotaxy across most families (except for Bromeliaceae, Mayacaceae, Cannaceae, Costaceae, Musaceae, and scattered cases in other families, e.g., *Palisota* Rchb. ex Endl. in Commelinaceae [Faden, 1988; Givnish et al., 1999]). The sister-group relationship of Arecales and Dasypogonales is supported by the possession of a woody habit in all of the former and most of the latter. Identification of woodiness as a synapomorphy for Arecales–Dasypogonales must await further evidence on relationships within Dasypogonaceae.

ANCESTRAL HABITAT RECONSTRUCTIONS

The ancestral habitat of Poales appears to have been sunny, damp to wet, highly infertile, and fireswept (see also Givnish et al., 1999; Linder & Rudall, 2005). Today, these conditions typify the family, either as a whole or in early divergent members (e.g., *Brocchinia*) in six cases: Bromeliaceae, Rapateaceae, Eriocaulaceae, Xyridaceae, Eceidocoleaceae, and Restionaceae (see Figs. 4, 5). However, the most species-rich families, Poaceae and Cyperaceae, often occur on more fertile substrates. Relatively few families and species are associated with forest understory environments (Fig. 4A), which helps to explain the high frequency of wind pollination within the order.

REPEATED EVOLUTION OF WIND POLLINATION IN POALES

Analyses based on our ML phylogeny indicate that wind pollination has arisen at least five times within Poales—in Typhaceae, the cyperids, the restiids, Eceidocoleaceae, and the PACCMAD-BEP clade of Poaceae (Fig. 6). This contrasts with three origins of wind pollination in Poales as inferred by Givnish et al. (1999) and Linder and Rudall (2005). Partly, this difference in conclusions is a result of different branching topologies, and partly, of different assumptions as to which taxa are wind pollinated. We assumed that insects pollinate *Flagellaria* and *Joinvillea*, based on arguments by Soreng and Davis (1998). We also assumed that insects pollinate *Anomochloa*, *Streptochaeta*, and *Pharus* P. Browne in Poaceae. Nothing certain is known about the pollination of these grasses. But the fact that all three lack feathery stigmas and versatile anthers on slender stamen filaments, both of which are usually associated with wind pollination, argues for their pollination by animals (Soderstrom, 1981; Soreng & Davis, 1998). In addition, *Anomochloa*, *Streptochaeta*, and at least one member of Pharoideae have grouped pollen granules with high exine relief, characters that are not associated with wind pollination in Poaceae (Page, 1978). Insect pollination has also been observed or inferred in a number of forest-dwelling bamboo grasses in herbaceous tribes Olyreae, Parianeae, and woody Bambuseae (Soderstrom & Calderón, 1971, 1979; Chapman, 1990; Salgado-Labouriau et al., 1992; Soreng & Davis, 1998); the placement of bamboos in the BEP clade (Grass Phylogeny Working Group I, 2001; Bouchenak-Khelladi et al., 2008) suggests this represents at least one secondary origin of animal pollination within the PACCMAD-BEP clade. The phyletic gap between the wind-pollinated graminids and the restiids of open habitats—involving animal-pollinated *Flagellaria* and *Joinvillea* in rain-forest understories and light gaps—clearly argues that any additional case of wind pollination in the graminids would be an additional independent origin. In addition, animal pollination in Anomochloideae and Pharoideae supports separate origins of wind pollination in Eceidocoleaceae and the PACCMAD-BEP clade of Poaceae based on our phylogeny (Fig. 6).

As predicted by several investigators and demonstrated across a wide sampling of angiosperms by Friedman and Barrett (2008), we found that animal pollination showed a significant correlation with floral showiness, flower size, floral sexuality, nectar production, and habitat openness (Table 1). Wind pollination was strongly associated, as predicted, with smaller, less conspicuous, often unisexual flowers, absence of nectar, and open habitats. Smaller and less

brightly colored flowers, often involving the loss of petals and/or sepals and nectar, are expected in wind-pollinated species given the energetic costs of these attractive structures, as well as the lack of utility of such structures in wind-pollinated taxa, and the possibility that the presence of petals and/or sepals, especially large ones, might interfere with the arrival of wind-borne pollen to stigmatic surfaces (see Linder, 1998; Culley et al., 2002; Linder & Rudall, 2005; Friedman & Barrett, 2008). It should be recognized that there is no need to invoke reevolution of petals and/or sepals in *Flagellaria* and *Joinvillea*, given the continuity of an ancestral line pollinated by animals up to the PACCMAD-BEP clade (Fig. 6).

Wind pollination and unisexual flowers—and their inverse, animal pollination with hermaphroditic flowers—should be associated because the reduced cost and increased benefits of attractive structures for attracting and dispersing pollen in hermaphroditic flowers applies only to animal-pollinated taxa (Givnish, 1980; Lloyd, 1982), and because unisexuality could help reduce self-pollination and stigma clogging in wind-pollinated taxa, given that the latter produce large amounts of pollen (Givnish, 1980; Lloyd & Webb, 1986; Charlesworth, 1993). The exceptional statistical significance of correlated evolution between wind pollination and unisexual flowers in analyses that include or exclude optimization of κ ($P < 0.0005$ and $P < 0.0002$, respectively) most likely has to do with the widespread occurrence of unisexual flowers in Poales, in nine of 16 families and a total of at least six clades.

Habitats that are open and windswept—even if only seasonally, as in the canopy of temperate deciduous forests—clearly favor wind pollination; conversely, windless understories favor animal pollination (Linder, 1998; Culley et al., 2002; Friedman & Barrett, 2008). However, the large number of animal-pollinated families of Poales found ancestrally in open habitats (Bromeliaceae, Eriocaulaceae, Mayacaceae, Xyridaceae) reduces the strength of the evolutionary correlation between habitat and pollination mechanism in the order. There is, however, a nearly perfect association of wind pollination with open habitats, including Typhaceae and the cyperids in open wetlands, and restiids, Ectepicolaceae, and the PACCMAD-BEP clade of Poaceae in open, often seasonally arid or fireswept upland habitats. Note the association of putative insect pollination with tropical forest understories and edges in *Flagellaria*, *Joinvillea*, and anomochloid, pharoid, and olyroid grasses (see Soderstrom & Calderón, 1971, and preceding discussion). Within-family reversions to insect pollination (not shown in Fig. 6) have occurred in scattered species in several genera of Cyperaceae

with brightly colored or fragrant inflorescences (e.g., *Ascolepis* Nees ex Steud., *Carex*, *Eleocharis* R. Br., *Hypolytrum* Rich., *Mapania* Aubl., *Rhynchospora* Vahl sect. *Dichromena* (Michx.) Griseb. [Goetghebeur, 1998; Magalhães et al., 2009]); several of these taxa grow in forest understories.

Nectar production should be negatively associated with wind pollination given the costs of producing nectar and its lack of utility in anemophilous species. Conversely, in some (but not all) animal-pollinated species, nectar serves as an attractant. The correlation between nectar production and pollination mechanism in Poales is significant but surprisingly weak (Table 1), with the latter probably reflecting the near absence of nectaries in the order outside the bromeliads. Even though Friedman and Barrett (2008) found that nectar production had the strongest pattern of correlated evolution with pollination mechanism of any factor they surveyed across the angiosperms, it would be difficult for such a pattern to occur in any lineage that is nearly uniform in nectar production or its absence. We coded *Potarophytum* Sandwith, the placeholder for Rapateaceae, as having septal nectaries, based on their presence in *Spathanthus* Desv. (Venturelli & Bouman, 1988) of tribe Rapateae and in *Guacamaya* Maguire, *Kunhardtia* Maguire, and *Schoenocephalum* Seub. of tribe Schoenocephalieae (Givnish et al., 2004). However, such nectaries are probably only of sporadic occurrence in Rapateaceae; they have not been observed as yet in other members of the family, and Renner (1989) described buzz pollination for *Saxo-fridericia* R. H. Schomb. (tribe Saxofridericieae) and *Stegolepis* Klotzsch ex Körn. (tribe Stegolepideae). If we assume that the ancestral condition for Rapateaceae is a lack of nectaries, then the correlation between pollination mechanism and nectar production remains weakly significant ($P < 0.045$ for $\kappa = 1$ and $P < 0.032$ for optimal κ).

Nectaries appear to have evolved independently in Eriocaulaceae, variously associated with petal appendages, staminodes, pistillodes, and pistils (Stützel, 1986; Rosa & Scatena, 2003, 2007; Ramos et al., 2005; Oriani et al., 2009), and in Xyridaceae, associated with stylar appendages in *Abolboda* (Stützel, 1990). *Flagellaria* secretes large amounts of nectar, with high concentrations of sugar and amino acids, from extrafloral nectaries and attracts large numbers of ants (Blüthgen et al., 2004a, b; Blüthgen & Fiedler, 2004); the ants observed to visit the flowers (Newell, 1969) almost surely are attracted by these extrafloral nectaries. Scoring *Flagellaria* as lacking nectar production (it has no floral nectaries) has almost no effect on the significance of correlated evolution between pollination mechanism and nectar

production, with $P < 0.047$ for tests with and without optimization of κ . If scoring were reversed for both *Flagellaria* and Rapateaceae, the correlation between pollination mechanism and nectar production would be marginally nonsignificant ($P < 0.058$ for $\kappa = 1$ and $P < 0.064$ for optimal κ).

The near absence of dioecy in Poales outside the restiids probably explains the nonsignificance of correlated evolution between wind pollination and dioecy in the order (Table 1). Although such a pattern is manifest across angiosperms, only the restiids show a high proportion of dioecious species within Poales, with scattered dioecious species or populations—unlikely to be scored in an analysis involving so few terminals as the current study—in Cyperaceae, Eriocaulaceae, Poaceae, and Typhaceae (Connor, 1981; Linder & Rudall, 2005). Low levels of dioecy at the familial level in Poales might be expected as based on the near absence of the woody habit and fleshy fruits at that level, given that both traits are strongly associated with dioecy across angiosperms and gymnosperms (Givnish, 1980; Renner & Ricklefs, 1995).

Finally, phylogenetic conservatism appears to account for the lack of a significant correlation between wind pollination and ovule number. Reduction to a single ovule is expected in wind-pollinated taxa given the presumed low probability of multiple pollen grains arriving on a single stigma (Dowding, 1987; Linder, 1998; Friedman & Barrett, 2008; but see Friedman & Barrett, 2009). However, in Poales, single ovules per carpel characterize the entire restiid–graminid clade irrespective of pollination mode, while multiple ovules characterize many of the taxa in the grade including Rapateaceae, the cyperids, and xyrids, again largely independent of pollination mode.

Two general questions that have not been satisfactorily addressed by previous authors remain: (1) why is wind pollination so widespread in Poales? and (2) why has wind pollination evolved five times independently in a single monocot order, when wind pollination is so rare in monocots? Of the factors just tested for correlated evolution with wind pollination, only one—open habitats—could be considered a prime mover, an external set of conditions that would drive the evolution of wind pollination and the many traits associated with the syndrome, namely, unisexual flowers, lack of nectar, reduced perianth, feathery stigma, versatile anthers, reduction in carpel and ovule number, and abundant, smooth, ulcerate pollen (Faegri & van der Pijl, 1979; Dahlgren & Clifford, 1982; Dahlgren et al., 1985; Linder, 1998; Friedman & Barrett, 2009). Potential factors favoring wind pollination identified by previous authors include

open, windswept habitats (see above), local dominance by conspecifics (Regal, 1982), and the absence or inefficiency of animal pollinators or the low quality of pollen delivered by them (Whitehead, 1983; Berry & Calvo, 1989; Cox, 1991; Weller et al., 1998; Culley et al., 2002). Strongly windswept habitats might also have poor animal visitation, so the first and third of these mechanisms are partly linked. Regal (1982) argued that high densities of conspecifics would favor wind-pollinated species, given the inherent inefficiencies of pollen delivery to conspecific stigmas in anemophily versus the higher costs of attractive structures or secretions in animal pollination. Indeed, several studies show that pollen delivery to conspecific stigmas drops rapidly with decreasing conspecific density in wind-pollinated taxa (Friedman & Barrett, 2009), and that such taxa have very high pollen:ovule ratios (Cruden, 1977, 2000).

But why have Poales evolved and retained wind pollination so frequently? One factor surely is that many early divergent lineages occupied habitats kept open by flooding, fire, and/or extreme soil infertility, and that open habitats were reinvaded by two additional lineages, Eceidicoleaceae and the PACC-MAD-BEP clade of Poaceae (Figs. 4–6). In addition, we propose that four additional prime movers favor wind pollination in herbs, including (1) tall stature, (2) vigorous vegetative spread, (3) adaptation to patchy disturbance by fire and/or flooding, and (4) positive feedback on conspecific abundance. These traits would all help generate high local dominance—and (1) through (3) above characterize all lineages that evolved or retained wind pollination in Poales, except for Eceidicoleaceae and Centrolepidaceae. Wind pollination in the latter two families may have been favored selection for reproductive assurance in the harsh, windswept alpine habitats and extremely infertile, fireswept, or seasonally inundated microsites they occupy. Goodwillie (1999) argued that wind pollination could provide such assurance—without the disadvantage of inbreeding associated with self-pollination—in open, harsh environments where pollinator abundance can vary greatly through time and space. Such assurance may be especially important for annual plants, given their need to produce seeds at the end of each growing season if an individual's genes are to enter the next generation, and given the likely spatial autocorrelation of poor conditions for pollinators.

Tall stature, vigorous vegetative spread, adaptation to patchy disturbances, and/or positive feedback on conspecific abundance—coupled with occupancy of open habitats—appear to provide a logical explanation for the distribution of wind pollination in other members of Poales. Typhaceae grow in open, fertile

habitats subject to frequent flooding and siltation, which are inimical to seedling establishment in most species, and to which few other species are adapted; Typhaceae gain high local dominance through advantages in height, rapid rhizomatous spread, and their ability to grow in wet, anoxic soils. Many bromeliads and rapateads lack rhizomatous growth or grow below closed canopies, where wind pollination should be a disadvantage. However, there are some genera (e.g., *Brocchinia* [Bromeliaceae] and *Stegolepis* [Rapateaceae]) in these two families that spread and form extensive, dense colonies on wet soil or rock surfaces, and it is not clear why wind pollination has not evolved in them. Extremely low growth rates in such habitats might prevent such plants from spreading and reaching local dominance for a substantial period after initial colonization of area, working against wind pollination. The origin of wind pollination in the cyperids may be related to invasion and rapid rhizomatous spread within frequently flooded or burnt areas. *Thurnia* of Thurniaceae forms massive colonies in streams on the Guayana Shield where few other plants grow; *Pronium* of the same family forms large monocultures in seasonal streambeds on sandstone in South Africa, which are flooded and burnt at frequent intervals.

Retention of animal pollination by Eriocaulaceae, Xyridaceae, and Mayacaceae despite their occurrence in frequently flooded or burnt sites may be related to their short stature and limited ability to achieve local dominance and spread laterally for any substantial distance. Their small size and slow growth almost surely reflect their occurrence in areas with wet, exceedingly poor soil. They usually inhabit sandy seepage zones and seasonal ponds where nutrients are not delivered in substantial amounts of transported sediment, unlike large-statured *Thurnia* and *Pronium* found in streams on similar sand or sandstone substrates.

Invasion of fire-prone, summer-dry upland habitats over infertile soils, as well as seasonally flooded microsites, by the restiids may help account for their local dominance and evolution of wind pollination (Linder & Rudall, 2005). Frequent, heavy rainfall and strong winds in some parts of western and montane Tasmania—where several early divergent taxa occur—might also favor wind pollination in restiids for pollination assurance. We argue (see above) that pollinator assurance was the primary driver toward anemophily in annual Centrolepidaceae that inhabit extremely infertile, seasonally wet depressions and rock pockets in southern Australia (Pignatti & Pignatti, 1994, 2005; Cooke, 2010). Pollination assurance probably also drove the origin and maintenance of wind pollination in tiny, perennial centrolepid cushion plants at high elevations in Tasmania and New Guinea, given

the harsh conditions for pollinators there and the extremely short stature and lack of local dominance of the centrolepids. Interestingly, both kinds of communities are also inhabited by strikingly convergent species of *Trithuria* (Nymphaeales: Hydatellaceae). Preliminary studies suggest that at least one of these (*T. submersa* Hook. f., in Western Australia) is also wind pollinated (Taylor & Williams, 2009).

Reinvasion of shaded habitats on more fertile, well-drained, unburnt sites by *Flagellaria* and *Joinvillea* favored reevolution of animal pollination, accompanied in *Flagellaria* by a small but conspicuously white corolla and scented flowers (Backer, 1951), abundant supply of extrafloral nectar (Blüthgen et al., 2004a, b; Blüthgen & Fiedler, 2004), and visitation by ants (Newell, 1969). Apparent retention of animal pollination in the anomoochloid and pharoid grasses presumably was favored by their restriction to forest understories. Finally, evolution and retention of wind pollination in the vast PACCMAD-BEP clade of Poaceae appear to have been favored by invasion of open habitats and achievement of high local dominance. Factors promoting such dominance might include (1) moderate to tall stature, usually combined with rapid rhizomatous spread; (2) morphological and physiological adaptations of grasses (e.g., narrow, erect foliage, C₄ photosynthesis) to widespread drought under bright, warm conditions starting in the Eocene and intensifying in the Miocene (Kellogg, 2000; Edwards & Smith, 2010); (3) positive feedback among grasses, fire, and nitrogen, given the low nitrogen content and decomposition rate of C₄ grasses and thus their flammability, the tolerance of grasses to fire due to their basal leaf meristems, the volatilization of nitrogen during fires, and the low nitrogen requirement of C₄ grasses (Seastedt et al., 1991; Wedin, 1995; Blair, 1997; Knapp et al., 1998; Reed et al., 2005); (4) positive feedback between grasses and grazers, given the attractiveness of grasses to many grazers, their resistance to grazing damage conferred by basal meristems, and collateral damage by grazers to other plants while seeking grasses (Archibald, 2008); and (5) positive feedback between grasses and their horizontally and vertically transmitted fungal endophytes (e.g., *Epichloë* Tul. & C. Tul., Clavicipitaceae, Ascomycota), based on protection against herbivores provided by the alkaloids secreted by the endophytes, resulting increases in the local frequency of infected grasses, and consequent increases in transmission of the endophytes to uninfected grasses (see Rudgers et al., 2009). An additional factor favoring at least temporary local dominance in grasses may involve (6) the negative effect that certain grasses have on nongrasses, at least in the short term, via the very large absorptive root surface developed by

grasses and their consequent ability to reduce the levels of soil nitrate available to other plants (see Craine, 2006; Dybzinski & Tilman, 2007). The great dominance of pooid grasses in cool areas (Edwards & Smith, 2010) might reflect (1), (4), (5), and/or (6), or some as yet unidentified ecological advantage of that group; the dominance of panicoid grasses in warmer, dryer areas probably reflects at least factors (1) through (4). Clearly, however, none of the proposed factors would have operated in early divergent grasses native to tropical forest understories. Wind pollination in most woody bamboos probably reflects its origin in the PACCMAD-BEP clade (Fig. 6). The alternative—that it represents yet another gain (i.e., a transition from animal-pollinated bamboos)—seems unlikely, given the restriction of known instances of animal pollination to a few understory genera of herbaceous bamboos, such as *Eremitis* Döll, *Olyra* L., *Pariana* Aubl., and *Piresta* Swallen (Gagné, 1969; Soderstrom & Calderón, 1971, 1979), which appear to be sister to the woody bamboos (see Bouchenak-Khelladi et al., 2008). Animal pollination might occur in a few woody bamboos (e.g., *Chusquea* Kunth; Janzen, 1976). If it does, it might represent a secondary gain of animal pollination, but our knowledge of bamboo pollination is far too rudimentary to hazard any analysis at this time. The retention of wind pollination in woody bamboos might be due to (7) their inability to sustain animal pollinators over the multi-year intervals between mass flowering events, driven by selection to satiate seed predators that feed on the unusually numerous (and sometimes unusually large) bamboo seeds, which like other grass seeds are nutritious and chemically unprotected (Janzen, 1976).

Our reanalysis of the evolution of wind pollination in Poales has thus produced several new insights, including a revision upward from three to five independent origins of wind pollination. In addition, we have shown that some traits that show a significant correlation with wind pollination across the angiosperms fail to do so within Poales, apparently due to phylogenetic conservatism in ovule number and non-dioecious breeding systems. Our analyses confirm that open habitats, lack of nectar production, and small, nonshowy, unisexual flowers show significant patterns of correlated evolution with wind pollination. Finally, we have proposed specific mechanisms—including several new ideas regarding the potential significance of plant stature, vegetative spread, and local positive feedback—to account for each of the five origins of wind pollination in Poales. Our predictions call for several new tests and for rigorous studies of pollination biology in some critical taxa (e.g., *Flagellaria*, *Joinvillea*, and the anomochlooid, pharoid, puelioid, and bambusoid grasses). Lynn Clark (pers. comm.) suggests

that pollination studies on relatively widespread *Pharus* might be especially interesting.

CONCLUSION

Deriving a monocot phylogeny based on plastomes—a core goal of the Monocot AToL Team—is designed to increase, to the maximum extent possible, the amount of phylogenetic information that can be wrung from the plastid genome. Sampling ever-greater numbers of bases per taxon should increase the resolution and likely accuracy of the resulting estimates of phylogenetic relationships in large taxonomic groups when combined with a reasonably dense and well-stratified sampling of taxa (Hillis, 1996; Givnish & Sytsma, 1997; Graybeal, 1998; Soltis et al., 1998; Hillis et al., 2003; Graham et al., 2006; Jansen et al., 2007). Our plastid data, including 25,107 informative sites, represent a 38-fold increase in the number of such sites over the pioneering study of monocot phylogeny by Duvall et al. (1993b), based on sequences of the single plastid gene *rbcL* and a comparable number of taxa. In the intervening years, monocot researchers increased the number of informative characters by sequencing genes that evolve faster (e.g., Fuse & Tamura, 2000, for *matK*; Givnish et al., 2005, for *ndhF*) or by concatenating and analyzing the sequences of several genes. The latter, multigene strategy has proven the more productive approach (e.g., see Chase et al., 2000; Soltis et al., 2000; Qiu et al., 2005), partly due to the difficulty of aligning rapidly evolving regions across highly divergent taxa, and to the inherently higher rate of homoplasy associated with such regions (Givnish & Sytsma, 1997). Chase et al. (2006) used 4777 informative sites from seven genes from the plastid, nuclear, and mitochondrial genomes to reconstruct relationships among 136 monocot taxa; Graham et al. (2006) used 5617 informative sites from 17 plastid genes and spacers to reconstruct ties among 94 monocot taxa. These studies increased the number of informative characters 5.3- to 8.6-fold over Duvall et al. (1993b) and increased the number of informative characters per taxon from 6.3 to 26 to 60.

The study presented here—using 81 plastid genes and over 100 kb of aligned sequence data per taxon—represents a further increase of fourfold to sixfold in data density per taxon, resulting in a ratio of 302 informative sites per taxon. Our study, and similar ones recently using 61 to 81 plastid gene sequences (Goremykin et al., 2003, 2004; Leebens-Mack et al., 2005; Cai et al., 2006; Jansen et al., 2007; Moore et al., 2007) to address angiosperm and land-plant relationships, mark the transition from plant multi-gene phylogenetics to phylogenomics. However, it

must be recognized that obtaining an avalanche of data—by sequencing plastomes in this project and, ultimately, transcriptomes and whole nuclear genomes in future analyses—will not relieve us of the need to complement large amounts of data per taxon with adequate taxon sampling. Long-branch attraction is a challenge that must constantly be kept in mind (Soltis & Soltis, 2004; Leebens-Mack et al., 2005; Whitfield & Lockhart, 2007). In the current study, given the more than fivefold variation in evolutionary rates in commelinids, the presence of numerous very long and very short branches in immediate proximity to each other, the conflict between the results of MP and ML analyses, and the current density of taxon sampling, long-branch attraction is clearly an issue. Relationships of the xyrids to each other, and to other members of the order, manifestly need additional taxon sampling and analysis to resolve. The appropriate level of taxon sampling density required to complement ever-rising amounts of sequence data per taxon is a central issue that must continue to be confronted theoretically (e.g., Geuten et al., 2007) and empirically (e.g., Soltis & Soltis, 2004; Leebens-Mack et al., 2005). To address these and other concerns, in the near future we plan to increase taxon sampling in Poales several-fold, and to explore additional analytical techniques (e.g., Lartillot et al., 2007) that may be less sensitive to long-branch attraction. Increased taxon sampling will also increase the power of comparative studies and of searches for morphological and anatomical synapomorphies for individual clades, and make careful calibration of our molecular tree and biogeographic analyses possible. The results presented here, however, show that even with the existing extent of taxon sampling, inferring a phylogeny based on whole plastid genomes can produce numerous new insights into monocot relationships and patterns of adaptive evolution at broad scales. The future of such studies—coupling a greatly increased number of plastomes with the sequencing of transcriptomes, the scoring and analysis of hundreds of morphological characters across representatives of all monocot families and subfamilies, the refinement of a timeline for monocot evolution, and analysis of broadscale patterns of monocot historical biogeography and adaptive evolution—appears very bright.

Literature Cited

- Angiosperm Phylogeny Group. 1998. An ordinal classification for the families of flowering plants. *Ann. Missouri Bot. Gard.* 85: 531–553.
- . 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Bot. J. Linn. Soc.* 141: 399–436.
- . 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot. J. Linn. Soc.* 161: 105–121.
- Archibald, S. 2008. African grazing lawns—How fire, rainfall, and grazer numbers interact to affect grass community states. *J. Wildlife Managem.* 72: 492–501.
- Asano, T., T. Tsudzuki, S. Takahashi, H. Shimada & K. Kadowski. 2004. Complete nucleotide sequence of the sugarcane (*Saccharum officinarum*) chloroplast genome: A comparative analysis of four monocot chloroplast genomes. *DNA Res.* 11: 93–99.
- Backer, C. A. 1951. Flagellariaceae. Pp. 245–250 in C. G. G. J. Van Steenis (editor), *Flora Malesiana*, Vol. 1. Noordhoff-Kolff, Jakarta.
- Baker, W. J., V. Savolainen, C. B. Asmussen-Lange, M. W. Chase, J. Dransfield, F. Forest, M. M. Harley, N. W. Uhl & M. Wilkinson. 2009. Complete generic-level phylogenetic analyses of palms (Arecaceae) with comparison of super-tree and supermatrix approaches. *Syst. Bot.* 58: 249–256.
- Bayer, C. & O. Appel. 1998. Joinvilleaceae. Pp. 249–251 in K. Kubitzki, H. Huber, P. J. Rudall, P. S. Stevens & T. Stützel (editors), *The Families and Genera of Vascular Plants*, Vol. 4. Flowering Plants, Monocotyledons: Alismatanae and Commelinanae (Except Gramineae). Springer Verlag, Berlin.
- Berry, P. E. & R. N. Calvo. 1989. Wind pollination, self-incompatibility, and altitudinal shifts in pollination systems in the high Andean genus *Espeletia* (Asteraceae). *Amer. J. Bot.* 76: 1602–1614.
- Blair, J. M. 1997. Fire, N availability, and plant response in grasslands: A test of the transient maxima hypothesis. *Ecology* 78: 2359–2368.
- Blüthgen, N. & K. Fiedler. 2004. Preferences for sugars and amino acids and their conditionality in a diverse nectar-feeding ant community. *J. Anim. Ecol.* 73: 155–166.
- , N. E. Stork & K. Fiedler. 2004a. Bottom-up control and co-occurrence in complex communities: Honeydew and nectar determine a rainforest ant mosaic. *Oikos* 106: 344–358.
- , G. Gottsberger & K. Fiedler. 2004b. Sugar and amino acid composition of ant-attended nectar and honeydew sources from an Australian rainforest. *Austral. Ecol.* 29: 418–429.
- Bosch, T., C. Löhne & J. Wiersema. 2008. Phylogeny and evolutionary patterns in Nymphaeales: Integrating genes, genomes and morphology. *Taxon* 57: 1052–1081.
- Bouchenak-Khelladi, Y., N. Salamin, V. Savolainen, F. Forest, M. van der Bank, M. W. Chase & T. R. Hodkinson. 2008. Large multi-gene phylogenetic trees of the grasses (Poaceae): Progress towards complete tribal and generic level sampling. *Molec. Phylog. Evol.* 47: 488–505.
- Bremer, K. 2000. Early Cretaceous lineages of monocot flowering plants. *Proc. Natl. Acad. U.S.A.* 97: 4707–4711.
- . 2002. Gondwanan evolution of the grass alliance of families (Poales). *Evolution* 56: 1374–1387.
- Briggs, B. G. & L. A. S. Johnson. 1998. *Georgeantha hexandra*, a new genus and species of Ecdiocoleaceae (Poales) from Western Australia. *Telopea* 7: 307–312.
- & H. P. Linder. 2009. A new subfamilial and tribal classification of Restionaceae (Poales). *Telopea* 12: 333–345.
- , A. D. Marchant, S. Gilmore & C. L. Porter. 2000. A molecular phylogeny of Restionaceae and allies. Pp. 661–671 in K. L. Wilson & D. Morrison (editors), *Monocots—Systematics and Evolution*. CSIRO, Melbourne.

- , ——— & A. J. Perkins. 2010. Phylogeny and features in Restionaceae, Centrolepidaceae and Anarthriaceae (the restioid clade of Poales). Pp. 357–388 in O. Seberg, G. Petersen, A. Barford & J. I. Davis (editors), *Diversity, Phylogeny, and Evolution in the Monocotyledons*. Aarhus University Press, Aarhus.
- Cai, Z. Q., C. Penaflor, J. V. Kuehl, J. Leebens-Mack, J. E. Carlson, C. W. dePamphilis, J. L. Boore & R. K. Jansen. 2006. Complete plastid genome sequences of *Drimys*, *Liriodendron*, and *Piper*: Implications for the phylogenetic relationships of magnoliids. *BMC Evol. Biol.* 6: 77.
- Cameron, K. M. 2004. Utility of plastid *psaB* gene sequences for investigating intrafamilial relationships within Orchidaceae. *Molec. Phylog. Evol.* 31: 1157–1180.
- , M. W. Chase & P. J. Rudall. 2003. Recircumscription of the monocotyledonous family Petrosaviaceae to include *Japonolirion*. *Brittonia* 55: 214–225.
- Campbell, C. S. & E. A. Kellogg. 1987. Sister group relationships of the Poaceae. Pp. 217–224 in T. R. Soderstrom, K. W. Hilu, C. S. Campbell & M. E. Barkworth (editors), *Grass Systematics and Evolution*. Smithsonian Institution, Washington, D.C.
- Chang, C. C., H. C. Lin, I. P. Lin, T. Y. Chow, H. H. Chen, W. H. Chen, C. H. Cheng, C. Y. Lin, S. M. Liu, C. C. Chang & S. M. Chaw. 2006. The chloroplast genome of *Phalaenopsis aphrodite* (Orchidaceae): Comparative analysis of evolutionary rate with that of grasses and its phylogenetic implications. *Molec. Biol. Evol.* 23: 279–291.
- Chang, J. T. 1996. Inconsistency of evolutionary tree reconstruction techniques when substitution rates vary across characters. *Math Biosci.* 134: 189–215.
- Chapman, G. P. 1990. The widening perspective: Reproductive biology of bamboos, some dryland grasses and cereals. Pp. 240–257 in G. P. Chapman (editor), *Reproductive Versatility in the Grasses*. Cambridge University Press, Cambridge.
- Charlesworth, D. 1993. Why are unisexual flowers associated with wind pollination and unspecialized pollinators? *Amer. Naturalist* 141: 481–490.
- Chase, M. W. 2004. Monocot relationships: An overview. *Amer. J. Bot.* 91: 1645–1655.
- , D. E. Soltis, R. G. Olmstead, D. Morgan, D. H. Les, B. D. Mishler, M. R. Duvall, R. Price, H. G. Hills, Y.-L. Qui, K. A. Kron, J. H. Rettig, E. Conti, J. D. Palmer, J. R. Manhart, K. J. Sytsma, H. J. Michaels, W. J. Kress, K. G. Karol, W. D. Clark, M. Hedren, B. S. Gaut, R. K. Jansen, K.-J. Kim, C. F. Wimpee, J. F. Smith, G. R. Furnier, S. H. Strauss, Q.-Y. Xiang, G. M. Plunkett, P. S. Soltis, S. E. Williams, P. A. Gadek, C. J. Quinn, L. E. Eguarte, E. Golenberg, G. H. Learn, S. Graham, S. C. H. Barrett, S. Dayanandan & V. A. Albert. 1993. Phylogenetics of seed plants: An analysis of nucleotide sequences from the plastid gene *rbcL*. *Ann. Missouri Bot. Gard.* 80: 528–580.
- , M. R. Duval, H. G. Hillis, J. G. Conran, A. V. Cox, L. E. Eguarte, J. Hartwell, M. F. Fay, L. R. Caddick, K. M. Cameron & S. Hoot. 1995a. Molecular phylogenetics of Liliales. Pp. 109–137 in P. J. Rudall, P. J. Cribb, D. F. Cutler & C. J. Humphries (editors), *Monocotyledons: Systematics and Evolution*. Royal Botanic Gardens, Kew, Richmond.
- , D. W. Stevenson, P. Wilkin & P. J. Rudall. 1995b. Monocot systematics: A combined analysis. Pp. 685–730 in P. J. Rudall, P. J. Cribb, D. F. Cutler & C. J. Humphries (editors), *Monocotyledons: Systematics and Evolution*. Royal Botanic Gardens, Kew, Richmond.
- , D. E. Soltis, P. S. Soltis, P. J. Rudall, M. F. Fay, W. J. Hahn, S. Sullivan, J. Joseph, M. Molvray, P. J. Kores, T. J. Givnish, K. J. Sytsma & J. C. Pires. 2000. Higher-level systematics of the monocotyledons: An assessment of current knowledge and a new classification. Pp. 3–16 in K. L. Wilson & D. A. Morrison (editors), *Monocots: Systematics and Evolution*. CSIRO Publishing, Collingwood, Victoria, Australia.
- , M. F. Fay, D. Devey, N. Rønstad, J. Davies, Y. Pillon, G. Petersen, O. Seberg, C. B. Asmussen, K. Hilu, T. Borsch, J. I. Davis, D. W. Stevenson, J. C. Pires, T. J. Givnish, K. J. Sytsma & S. W. Graham. 2006. Multi-gene analyses of monocot relationships: A summary. Pp. 63–75 in J. T. Columbus, E. A. Friar, C. W. Hamilton, J. M. Porter, L. M. Prince & M. G. Simpson (editors), *Monocots: Comparative Biology and Evolution* (excluding Poales). Rancho Santa Ana Botanic Garden, Claremont, California.
- Chevreur, B., T. Pfisterer, B. Drescher, A. J. Driesel, W. E. G. Müller, T. Wetter & S. Suhai. 2004. Using the miraEST Assembler for reliable and automated mRNA transcript assembly and SNP Detection in sequenced ESTs. *Genome Res.* 14: 1147–1159.
- Christin, P. A., G. Besnard, E. Samaritani, M. R. Duvall, T. R. Hodkinson, V. Savolainen & N. Salamin. 2008. Oligocene CO₂ decline promoted C₄ photosynthesis. *Curr. Biol.* 18: 37–43.
- Clark, L. G., W. Zhang & J. F. Wendel. 1995. A phylogeny of the grass family (Poaceae) based on *ndhF* sequence data. *Syst. Bot.* 20: 436–460.
- , M. Kobayashi, S. Mathews, R. E. Spangler & E. A. Kellogg. 2000. The Puelioideae, a new subfamily of Poaceae. *Syst. Bot.* 25: 181–187.
- Clifford, H. T., G. J. Keighery & J. G. Conran. 1998. Dasygogonaceae. Pp. 190–194 in K. Kubitzki, H. Huber, P. J. Rudall, P. S. Stevens & T. Stützel (editors), *The Families and Genera of Vascular Plants, Vol. 4. Flowering Plants, Monocotyledons: Alismatanae and Commelinanae* (Except Gramineae). Springer Verlag, Berlin.
- Columbus, J. T., E. A. Friar, C. W. Hamilton, J. M. Porter, L. M. Prince & M. G. Simpson (editors). 2006. *Monocots: Comparative Biology and Evolution* (Excluding Poales). Rancho Santa Ana Botanic Garden, Claremont, California.
- , ———, ———, ———, ——— & ——— (editors). 2007. *Monocots: Comparative Biology and Evolution* (Poales). Rancho Santa Ana Botanic Garden, Claremont, California.
- Connor, H. E. 1981. Evolution of the reproductive systems in the Gramineae. *Ann. Missouri Bot. Gard.* 68: 48–74.
- Conran, J. G., R. J. Carpenter & G. J. Jordan. 2009. Early Eocene *Ripogonum* (Liliales: Ripogonaceae) leaf macrofossils from southern Australia. *Austral. Syst. Bot.* 22: 219–228.
- Cooke, D. A. 1998. Centrolepidaceae. Pp. 106–109 in K. Kubitzki, H. Huber, P. J. Rudall, P. S. Stevens & T. Stützel (editors), *The Families and Genera of Vascular Plants, Vol. 4. Flowering Plants, Monocotyledons: Alismatanae and Commelinanae* (Except Gramineae). Springer Verlag, Berlin.
- . 2010. A flora treatment of Australian Centrolepidaceae. <<http://members.ozemail.com.au/~davcooke/centrlep.htm>>, accessed 8 July 2010.
- Cox, A., A. Pridgeon, V. Albert & M. Chase. 1997. Phylogenetics of the slipper orchids (Cypripedioideae, Orchidaceae): Nuclear rDNA ITS sequences. *Pl. Syst. Evol.* 208: 197–223.
- Cox, P. A. 1991. Abiotic pollination: An evolutionary escape for animal-pollinated angiosperms. *Philos. Trans., Ser. B* 333: 217–224.

- Craine, J. M. 2006. Competition for nutrients and optimal root allocation. *Pl. & Soil* 235: 171–185.
- Cruden, R. W. 1977. Pollen-ovule ratios: A conservative indicator of breeding systems in flowering plants. *Evolution* 31: 32–46.
- . 2000. Pollen grains: Why so many? *Pl. Syst. Evol.* 222: 143–165.
- Culley, T. M., S. G. Weller & A. K. Sakai. 2002. The evolution of wind pollination in angiosperms. *Trends Ecol. Evol.* 17: 361–369.
- Dahlgren, R. M. T. & H. T. Clifford. 1982. *The Monocotyledons: A Comparative Study*. Academic Press, London.
- , ——— & P. F. Yeo. 1985. *The Families of the Monocotyledons*. Springer Verlag, Berlin.
- Davies, T. J., V. Savolainen, M. W. Chase, J. Moat & T. G. Barraclough. 2004. Environmental energy and evolutionary rates in flowering plants. *Proc. Roy. Soc. London, Ser. B, Biol. Sci.* 271: 2195–2200.
- Davis, J. L., D. W. Stevenson, G. Petersen, O. Seberg, L. M. Campbell, J. V. Freudenstein, D. H. Goldman, C. R. Hardy, F. A. Michelangeli, M. P. Simmons, C. D. Specht, F. Vergara-Silva & M. Gandolfo. 2004. A phylogeny of the monocots, as inferred from *rbcL* and *atpA* sequence variation, and a comparison of methods for calculating jackknife and bootstrap values. *Syst. Bot.* 29: 467–510.
- Dowding, P. 1987. Wind pollination mechanisms and aerobiology. *Int. Rev. Cytol.* 107: 421–437.
- Doyle, J. D., J. I. Davis, R. J. Soreng, D. Garvin & M. J. Anderson. 1992. Chloroplast DNA inversions and the origin of the grass family (Poaceae). *Proc. Natl. Acad. Sci. U.S.A.* 89: 7722–7726.
- Drábková, L. Z. & C. Vlček. 2009. DNA variation within Juncaceae: Comparison of impact of organelle regions on phylogeny. *Pl. Syst. Evol.* 278: 169–186.
- Dunn, R. R., A. D. Gove, T. G. Barraclough, T. J. Givnish & J. D. Majer. 2007. Convergent evolution of an ant-plant mutualism across plant families, continents, and time. *Evol. Ecol. Res.* 9: 1349–1362.
- Duvall, M. R., G. E. Learn Jr., L. E. Eguarte & M. T. Clegg. 1993a. Phylogenetic analysis of *rbcL* sequences identifies *Acorus calamus* as the primal extant monocotyledon. *Proc. Natl. Acad. Sci. U.S.A.* 90: 4641–4644.
- , M. T. Clegg, M. W. Chase, W. D. Clark, W. J. Kress, H. G. Hills, L. E. Eguarte, J. F. Smith, B. S. Gaut, E. A. Zimmer & G. H. Learn. 1993b. Phylogenetic hypotheses for the monocotyledons constructed from *rbcL* sequence data. *Ann. Missouri Bot. Gard.* 80: 607–619.
- Dybzinski, R. & D. Tilman. 2007. Resource use patterns predict long-term outcomes of plant competition for nutrients and light. *Amer. Naturalist* 170: 305–318.
- Edgar, R. C. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucl. Acids Res.* 32: 1792–1797.
- Edwards, E. J. & S. J. Smith. 2010. Phylogenetic analyses reveal the shady history of C_4 grasses. *Proc. Natl. Acad. Sci. U.S.A.* 107: 2532–2537.
- Ervik, F. 1993. Notes on the phenology and pollination of the dioecious palms *Mauritia flexuosa* and *Aphandra natalia* in Ecuador. Pp. 7–12 in W. Barthlott, C. M. Naumann, K. Schmidt-Loske & K.-L. Schuchmann (editors), *Animal-plant Interactions in Tropical Environments*. Zoologisches Forschungsinstitut und Museum Alexander König, Bonn.
- Faden, R. B. 1988. Vegetative and reproductive features of forest and nonforest genera of African Commelinaceae. Pp. 521–526 in P. Goldblatt & P. P. Lowry II (editors), *Modern Systematic Studies in African Botany*. Monogr. Syst. Bot. Missouri Bot. Gard. 25.
- Faegri, K. & L. van der Pijl. 1979. *The Principles of Pollination Ecology*, 3rd ed. Oxford University Press, Oxford.
- Felsenstein, J. 1985. Phylogenies and the comparative method. *Amer. Naturalist* 125: 1–15.
- Friedman, J. & S. C. H. Barrett. 2008. A phylogenetic analysis of the evolution of wind pollination in the angiosperms. *Int. J. Pl. Sci.* 169: 49–58.
- & ———. 2009. Winds of change: New insights on the ecology and evolution of pollination and mating in wind-pollinated plants. *Ann. Bot. (Oxford)* 103: 1515–1527.
- Friis, E. M., K. R. Pedersen & P. R. Crane. 2004. Araceae from the Early Cretaceous of Portugal: Evidence on the emergence of monocotyledons. *Proc. Natl. Acad. Sci. U.S.A.* 101: 16565–16570.
- Fuse, S. & M. N. Tamura. 2000. A phylogenetic analysis of the plastid *matK* gene with emphasis on Melanthiaceae. *Pl. Biol. (Stuttgart)* 2: 415–427.
- Gadagkar, S. R. & S. Kumar. 2005. Maximum likelihood outperforms maximum parsimony even when evolutionary rates are heterotachous. *Molec. Biol. Evol.* 22: 2139–2141.
- Gagné, R. J. 1969. A new genus and two new species of Cecidomyiidae associated with *Pariaria* spp. (Graminae) in South America (Diptera). *Proc. Entomol. Soc. Wash.* 71: 108–111.
- Gaut, B. S., S. V. Muse, W. D. Clark & M. T. Clegg. 1992. Relative rates of nucleotide substitution in the *rbcL* locus in monocotyledonous plants. *J. Molec. Evol.* 35: 292–303.
- , B. R. Morton, B. C. McCaig & M. T. Clegg. 1996. Substitution rates in grasses and palms: Synonymous rate differences at the nuclear gene *Adh* parallel rate difference in the plastid gene *rbcL*. *Proc. Natl. Acad. Sci. U.S.A.* 93: 10274–10279.
- Geuten, K., T. Massingham, P. Darius, E. Smets & N. Goldman. 2007. Experimental design criteria in phylogenetics: Where to add taxa. *Syst. Biol.* 56: 609–622.
- Givnish, T. J. 1980. Ecological constraints on the evolution of breeding systems—Dioecy and dispersal in gymnosperms. *Evolution* 34: 959–972.
- & K. J. Sytsma. 1997. Consistency, characters, and the likelihood of correct phylogenetic inference. *Molec. Phylogen. Evol.* 7: 320–330.
- , ———, J. F. Smith, W. J. Hahn, D. H. Benzing & E. M. Burkhardt. 1997. Molecular evolution and adaptive radiation in *Brocchinia* (Bromeliaceae: Pitcairnioideae) atop tepuis of the Guayana Shield. Pp. 259–311 in T. J. Givnish & K. J. Sytsma (editors), *Molecular Evolution and Adaptive Radiation*. Cambridge University Press, New York.
- , T. M. Evans, J. C. Pires & K. J. Sytsma. 1999. Polyphyly and convergent morphological evolution in Commelinales and Commelinidae: Evidence from *rbcL* sequence data. *Molec. Phylogen. Evol.* 12: 360–385.
- , ———, M. L. Zjhra, T. B. Patterson, P. E. Berry & K. J. Sytsma. 2000. Molecular evolution, adaptive radiation, and geographic diversification in the amphiatlantic family Rapateaceae: Evidence from *ndhF* sequences and morphology. *Evolution* 54: 1915–1937.
- , K. C. Millam, T. M. Evans, J. C. Hall, J. C. Pires, P. E. Berry & K. J. Sytsma. 2004. Ancient vicariance or recent long-distance dispersal? Inferences about phylogeny and South American–African disjunction in Rapateaceae and Bromeliaceae based on *ndhF* sequence data. *Int. J. Pl. Sci.* 165: S35–S54.
- , J. C. Pires, S. W. Graham, M. A. McPherson, L. M. Prince, T. B. Patterson, H. S. Rai, E. R. Roalson, T. M. Evans, W. J. Hahn, K. C. Millam, A. W. Meerow, M.

- McNeal, J. Leebens-Mack & L. Cui. 2005. Methods for obtaining and analyzing chloroplast genome sequences. *Meth. Enzymol.* 395: 348–384.
- , C. Caittani, C. Saski, S. B. Lee, J. Tomkins, A. J. Alverson & H. Daniell. 2006. Phylogenetic analyses of *Vitis* (Vitaceae) based on complete chloroplast genome sequences: Effects of taxon sampling and phylogenetic methods on resolving relationships among rosids. *BMC Evol. Biol.* 6: 32.
- , Z. Cai, L. A. Raubeson, H. Daniell, C. W. dePamphilis, J. Leebens-Mack, K. F. Müller, M. Guisinger-Bellian, R. C. Haberle, A. K. Hansen, T. W. Chumley, S.-B. Lee, R. Peery, J. R. McNeal, J. V. Kuehl & J. L. Boore. 2007. Analysis of 81 genes from 64 plastid genomes resolves relationships in angiosperms and identifies genome-scale evolutionary patterns. *Proc. Natl. Acad. Sci. U.S.A.* 104: 19369–19374.
- Janssen, T. & K. Bremer. 2004. The age of major monocot groups inferred from 800+ *rbcL* sequences. *Bot. J. Linn. Soc.* 146: 385–398.
- Janzen, D. H. 1976. Why bamboos wait so long to flower. *Ann. Rev. Ecol. Syst.* 7: 347–391.
- Johansen, L. B. 2005. Phylogeny of *Orchidantha* (Lowiaceae) and the Zingiberales based on six DNA regions. *Syst. Bot.* 30: 106–117.
- Jordan, G. 2009. Restionaceae. Key to Tasmanian vascular plants. <http://www.utas.edu.au/dicotkey/dicotkey/restio_etc/Restionaceae.htm>, accessed 1 July 2010.
- Kellogg, E. A. 2000. The grasses: A case study in macroevolution. *Ann. Rev. Ecol. Syst.* 31: 217–238.
- & H. P. Linder. 1995. Phylogeny of Poales. Pp. 511–542 in P. J. Rudall, P. J. Cribb, D. F. Cutler & C. J. Humphries (editors), *Monocotyledons: Systematics and Evolution*. Royal Botanic Gardens, Kew, Richmond.
- Kim, K. J. & H. L. Lee. 2004. Complete chloroplast genome sequences from Korean ginseng (*Panax schinseng* Nees) and comparative analysis of sequence evolution among 17 vascular plants. *DNA Res.* 11: 247–261.
- Knapp, A. K., J. M. Briggs, J. M. Blair & C. L. Turner. 1998. Patterns and controls of aboveground net primary production in tallgrass prairie. Pp. 193–221 in A. K. Knapp, J. M. Briggs, D. C. Hartnett & S. L. Collins (editors), *Grassland Dynamics: Long-term Ecological Research in Tallgrass Prairie*. Oxford University Press, New York.
- Kral, R. 1998. Xyridaceae. Pp. 461–469 in K. Kubitzki, H. Huber, P. J. Rudall, P. S. Stevens & T. Stützel (editors), *The Families and Genera of Vascular Plants, Vol. 4. Flowering Plants, Monocotyledons: Alismatanae and Commelinanae (Except Gramineae)*. Springer Verlag, Berlin.
- Kress, W. J., L. M. Prince, W. J. Hahn & E. A. Zimmer. 2001. Unraveling the evolutionary radiation of the families of the Zingiberales using morphological and molecular evidence. *Syst. Biol.* 50: 926–944.
- Lartillot, N., H. Brinkman & H. Philippe. 2007. Suppression of long-branch attraction artefacts in the animal phylogeny using a site-heterogeneous model. *BMC Evol. Biol.* 7: S4.
- Leebens-Mack, J., L. A. Raubeson, L. Cui, J. V. Kuehl, M. H. Fourcade, T. W. Chumley, J. L. Boore, R. K. Jansen & C. W. dePamphilis. 2005. Identifying the basal angiosperm node in chloroplast genome phylogenies: Sampling one's way out of the Felsenstein zone. *Molec. Biol. Evol.* 22: 1948–1963.
- Les, D. H., M. A. Cleland & M. Waycott. 1997. Phylogenetic studies in Alismatidae, II: Evolution of marine angiosperms (seagrasses) and hydrophily. *Syst. Bot.* 22: 443–463.
- Leseberg, C. H. & M. R. Duvall. 2009. The complete chloroplast genome of *Coix lachryma-jobi* and a comparative molecular evolutionary analysis of plastomes in cereals. *J. Molec. Evol.* 69: 311–318.
- Linder, H. P. 1987. The evolutionary history of the Poales/Restionales—A hypothesis. *Kew Bull.* 42: 297–318.
- . 1998. Morphology and the evolution of wind pollination. Pp. 123–125 in S. J. Owens & P. J. Rudall (editors), *Reproductive Biology in Systematics, Conservation and Economic Botany*. Royal Botanic Gardens, Kew, Richmond.
- & P. J. Rudall. 2005. Evolutionary history of the Poales. *Ann. Rev. Ecol. Evol. Syst.* 36: 107–124.
- , B. G. Briggs & L. A. S. Johnson. 1998. Ecdiocolaeaceae. Pp. 195–197 in K. Kubitzki, H. Huber, P. J. Rudall, P. S. Stevens & T. Stützel (editors), *The Families and Genera of Vascular Plants, Vol. 4. Flowering Plants, Monocotyledons: Alismatanae and Commelinanae (Except Gramineae)*. Springer Verlag, Berlin.
- , P. Eldenäs & B. G. Briggs. 2003. Contrasting patterns of radiation in African and Australian Restionaceae. *Evolution* 57: 2688–2702.
- Listabarth, C. 1992. Insect-induced wind pollination of the palm *Chamaedorea pinnatifrons* and pollination in the related *Wendlandiella* sp. *Biodivers. & Conservation* 1: 39–50.
- Lloyd, D. G. 1982. Selection of combined versus separate sexes in seed plants. *Amer. Naturalist* 120: 571–585.
- & C. J. Webb. 1986. The avoidance of interference between the presentation of pollen and stigmas in angiosperms. I. Dichogamy. *New Zealand J. Bot.* 24: 135–162.
- Maddison, D. R. 1991. The discovery and importance of multiple islands of most parsimonious trees. *Syst. Zool.* 40: 315–328.
- Maddison, W. P. & D. R. Maddison. 1992. *MacClade: Analysis of Phylogeny and Character Evolution (Vers. 3)*. Sinauer Associates, Sunderland, Massachusetts.
- Magalhães, A. F., A. L. T. G. Ruiz, A. Flach, A. D. Faria, E. G. Magalhães & M. do Carmo E. Amaral. 2009. Floral scent of *Eleocharis elegans* (Kunth) Roem. & Schult. (Cyperaceae). *Biochem. Syst. Ecol.* 33: 675–679.
- Maier, R. M., K. Neckermann, G. L. Iglói & H. Kossel. 1995. Complete sequence of the maize chloroplast genome—Gene content, hotspots of divergence and fine-tuning of genetic information by transcript editing. *J. Molec. Evol.* 25: 614–628.
- Marchant, A. D. & B. G. Briggs. 2007. Ecdiocolaeaceae and Joinvilleaceae, sisters of Poaceae (Poales): Evidence from *rbcL* and *matK* data. *Telopea* 11: 437–450.
- Mardanov, A. V., N. V. Ravin, B. B. Kuznetsov, T. H. Samigullin, A. S. Antonov, T. V. Kolganova & K. G. Skyabin. 2008. Complete sequence of the duckweed (*Lemna minor*) chloroplast genome: Structural organization and phylogenetic relationships to other angiosperms. *J. Molec. Evol.* 66: 555–564.
- Matsushima, R., Y. Hu, K. Toyoda, Sodmergen & W. Sakamoto. 2008. The model plant *Medicago truncatula* exhibits biparental plastid inheritance. *Pl. Cell Physiol.* 49: 81–91.
- Michelangeli, F. A., J. I. Davis & D. W. Stevenson. 2003. Phylogenetic relationships among Poaceae and related families as inferred from morphology, inversions in the plastid genome, and sequence data from the mitochondrial and plastid genomes. *Amer. J. Bot.* 90: 93–106.

- Moore, M. J., A. Dhingra, P. S. Soltis, R. Shaw, W. G. Farmerie, K. M. Foltá & D. E. Soltis. 2006. Rapid and accurate pyrosequencing of angiosperm plastid genomes. *BMC Pl. Biol.* 6: 17.
- , C. D. Bell, P. S. Soltis & D. E. Soltis. 2007. Using plastid genome-scale data to resolve enigmatic relationships among basal angiosperms. *Proc. Natl. Acad. Sci. U.S.A.* 104: 19363–19368.
- Morris, L. M. & M. R. Duvall. 2010. The chloroplast genome of *Anomochloa marantoidea* (Anomochloaceae; Poaceae) comprises a mixture of grass-like and unique features. *Amer. J. Bot.* 97: 620–627.
- Moura, J. I. L., F. J. Cividanes, L. P. dos Santos Filho & R. R. Valle. 2008. Pollination of oil palm by weevils in southern Bahia, Brazil. *Pesq. Agropecu. Brasil.* 43: 289–294.
- Muasya, A. M., D. A. Simpson, G. A. Verboom, P. Goetghebeur, R. F. C. Naczi, M. W. Chase & E. Smets. 2009. Phylogeny of Cyperaceae based on DNA sequence data: Current progress and future prospects. *Bot. Rev. (Lancaster)* 75: 2–21.
- Newell, T. K. 1969. A study of the genus *Joinvillea* (Flagellariaceae). *J. Arnold Arbor.* 50: 527–555.
- Ogihara, Y., K. Isono, T. Kojima, A. Endo, M. Hanaoka, T. Shiina, T. Terachi, S. Utsugi, M. Murata, N. Mori, S. Takumi, K. Ikeo, T. Gjobori, R. Murai, K. Murai, Y. Matsuoka, Y. Ohnishi, H. Tajiri & K. Tsunewaki. 2002. Structural features of a wheat plastome as revealed by complete sequencing of chloroplast DNA. *Molec. Genet. Genomics* 266: 740–746.
- Okumura, S., M. Sawada, Y. W. Park, T. Hayashi, M. Shimamura, H. Takase & K.-I. Tomizawa. 2006. Transformation of poplar (*Populus alba*) plastids and expression of foreign proteins in tree chloroplasts. *Transgen. Res.* 15: 637–646.
- Oriani, A., P. T. Sano & V. L. Scatena. 2009. Pollination biology of *Syngonanthus elegans* (Eriocaulaceae—Poales). *Austral. J. Bot.* 57: 94–105.
- Page, J. S. 1978. A scanning electron microscope survey of grass pollen. *Kew Bull.* 32: 313–319.
- Pagel, M. & A. Meade. 2006. Bayesian analysis of correlated evolution of discrete characters by reverse-jump Markov chain Monte Carlo. *Amer. Naturalist* 167: 808–825.
- Pignatti, E. & S. Pignatti. 1994. *Centrolepidi—Hydrocotyleta alatae*, a new class of ephemeral communities in Western Australia. *J. Veg. Sci.* 5: 55–62.
- & ———. 2005. Ephemeral wet vegetation in Western Australia. *Phytocoenologia* 35: 201–218.
- Pires, J. C., I. J. Maureira, T. J. Givnish, K. J. Sytsma, O. Seberg, G. Petersen, J. I. Davis, D. W. Stevenson, P. J. Rudall, M. F. Fay & M. W. Chase. 2006. Phylogeny, genome size, and chromosome evolution of Asparagales. Pp. 287–304 in J. T. Columbus, E. A. Friar, J. M. Porter, L. M. Prince & M. G. Simpson (editors), *Monocots: Comparative Biology and Evolution* Poales. Rancho Santa Ana Botanic Garden, Claremont, California.
- Plader, W., Y. Yukawa, M. Sugiura & S. Malepszy. 2007. The complete structure of the cucumber (*Cucumis sativus* L.) chloroplast genome: Its composition and comparative analysis. *Cell. Molec. Biol. Lett.* 12: 584–594.
- Qiu, Y. L., O. Dombrowska, J. Lee, B. A. Whitlock, F. Bernasconi-Quadroni, J. S. Rest, C. C. Davis, T. Borsch, K. W. Hilu, S. S. Renner, D. E. Soltis, P. S. Soltis, M. J. Zanis, J. J. Cannone, R. R. Gutell, M. Powell, V. Savolainen, L. W. Chatrou & M. W. Chase. 2005. Phylogenetic analyses of basal angiosperms based on nine plastid, mitochondrial, and nuclear genes. *Int. J. Pl. Sci.* 166: 815–842.
- Ramirez, S. R., B. Gravendeel, R. B. Singer, C. R. Marshall & N. E. Pierce. 2007. Dating the origin of the Orchidaceae from a fossil orchid with its pollinator. *Nature* 448: 1042–1045.
- Ramos, C. O. C., E. L. Borba & L. S. Funch. 2005. Pollination in Brazilian *Syngonanthus* (Eriocaulaceae) species: Evidence for entomophily instead of anemophily. *Ann. Bot. (Oxford)* 96: 387–397.
- Ratan, A. 2009. Assembly algorithms for next-generation sequence data. Ph.D. Dissertation, Department of Computer Science, Pennsylvania State University, State College, Pennsylvania.
- Raubeson, L. A., R. Peery, T. W. Chumley, C. Dziubek, H. M. Fourcade, J. L. Boore & R. K. Jansen. 2007. Comparative chloroplast genomics: Analyses including new sequences from the angiosperms *Nuphar advena* and *Ranunculus macranthus*. *BMC Genomics* 8: 174.
- Reed, H. E., T. R. Seastedt & J. M. Blair. 2005. Ecological consequences of C₄ grass invasion of a C₄ grassland: A dilemma for management. *Ecol. Applic.* 15: 1560–1569.
- Regal, P. J. 1982. Pollination by wind and animals: Ecology of geographic patterns. *Ann. Rev. Ecol. Syst.* 13: 497–524.
- Renner, S. S. 1989. Floral biological observations on *Heliamphora tatei* (Sarraceniaceae) and other plants from Cerro de la Neblina in Venezuela. *Pl. Syst. Evol.* 163: 21–29.
- & R. E. Ricklefs. 1995. Dioecy and its correlates in the flowering plants. *Amer. J. Bot.* 82: 596–606.
- Rosa, M. M. & V. L. Scatena. 2003. Floral anatomy of *Eriocaulon elichrysoides* and *Syngonanthus caulescens* (Eriocaulaceae). *Flora* 198: 188–199.
- & ———. 2007. Floral anatomy of Paepalanthoideae (Eriocaulaceae, Poales) and their nectariferous structures. *Ann. Bot. (Oxford)* 99: 131–139.
- Rudall, P. J. 1997. The nucellus and chalaza in monocotyledons: Structure and systematics. *Bot. Rev. (Lancaster)* 63: 140–181.
- & E. Caddick. 1994. Investigation of the presence of phenolic compounds in monocot cell walls, using UV fluorescence microscopy. *Ann. Bot. (Oxford)* 75: 483–491.
- , P. J. Cribb, D. F. Cutler & C. J. Humphries (editors). 1995. *Monocotyledons: Systematics and Evolution*, Vols. I and II. Royal Botanic Gardens, Kew, Richmond.
- , T. Eldridge, J. Tratt, M. M. Ramsay, R. E. Tuckett, S. Y. Smith, M. E. Collinson, M. V. Remizowa & D. D. Sokoloff. 2009. Seed fertilization, development, and germination in Hydatellaceae (Nymphaeales): Implications for endosperm evolution in early angiosperms. *Amer. J. Bot.* 96: 1581–1593.
- Rudgers, J. A., M. E. Afkhami, M. A. Rúa, A. J. Davitt, S. Hammer & V. M. Huguet. 2009. A fungus among us: Broad patterns of endophyte distribution in the grasses. *Ecology* 90: 1531–1539.
- Saarela, J. M. & S. W. Graham. 2010. Inference of phylogenetic relationships among the subfamilies of grasses (Poaceae: Poales) using meso-scale sampling of the plastid genome. *Botany* 88: 65–84.
- , H. S. Rai, J. A. Doyle, P. K. Endress, S. Mathews, A. D. Marchant, B. G. Briggs & S. W. Graham. 2007. Hydatellaceae identified as a new branch near the base of the angiosperm phylogenetic tree. *Nature* 446: 312–315.
- , P. J. Prentis, H. S. Rai & S. W. Graham. 2008. Phylogenetic relationships in the monocot order Commelinales, with a focus on Philydraceae. *Botany* 86: 719–731.
- Salgado-Labouriau, M. L., S. Nilsson & M. Rinaldi. 1992. Exine sculpture in *Pariana* pollen (Gramineae). *Grana* 32: 243–249.

- Samson, N., M. G. Bausher, S.-B. Lee, R. K. Jansen & H. Daniell. 2007. The complete nucleotide sequence of the coffee (*Coffea arabica* L.) chloroplast genome: Organization and implications for biotechnology and phylogenetic relationships amongst angiosperms. *Pl. Biotechnol. J.* 5: 339–353.
- Sánchez-Ken, J. G., L. G. Clark, E. A. Kellogg & E. A. Kay. 2007. Reinstatement and emendation of subfamily Micrairoideae. *Syst. Bot.* 32: 71–80.
- Saski, C., S. B. Lee, S. Fjellheim, C. Guda, R. K. Jansen, H. Luo, J. Tomkins, O. A. Rogdil, H. Daniell & J. L. Clarke. 2007. Complete chloroplast genome sequences of *Hordeum vulgare*, *Sorghum bicolor* and *Agrostis stolonifera*, and comparative analyses with other grass genomes. *Theor. Appl. Genet.* 115: 571–590.
- Schmitz-Linneweber, C., R. M. Maier, J. P. Alcaraz, A. Cottet, R. G. Herrmann & R. Mache. 2001. The plastid chromosome of spinach (*Spinacia oleracea*): Complete nucleotide sequence and gene organization. *Pl. Molec. Biol.* 45: 307–315.
- Seastedt, T. R., J. M. Briggs & D. J. Gibson. 1991. Controls of nitrogen limitation in tallgrass prairie. *Oecologia* 87: 72–79.
- Seberg, O., G. Petersen, A. S. Barford & J. I. Davis (editors). 2010. Diversity, Phylogeny, and Evolution in the Monocotyledons. Aarhus University Press, Aarhus.
- Simpson, D. A., C. A. Furness, T. R. Hodkinson, A. M. Muasya & M. W. Chase. 2003. Phylogenetic relationships in Cyperaceae subfamily Mapanioideae inferred from pollen and plastid DNA sequence data. *Amer. J. Bot.* 90: 1071–1086.
- Smith, S. A. & M. J. Donoghue. 2008. Rates of molecular evolution are linked to life history in flowering plants. *Science* 322: 86–89.
- Soderstrom, T. R. 1981. Some evolutionary trends in the Bambusoideae (Poaceae). *Ann. Missouri Bot. Gard.* 68: 15–47.
- & C. E. Calderón. 1971. Insect pollination in tropical rain forest grasses. *Biotropica* 3: 1–16.
- & ———. 1979. A commentary on the bamboos (Poaceae: Bambusoideae). *Biotropica* 11: 161–172.
- Sokoloff, D. D., M. V. Remizowa, H. P. Linder & P. J. Rudall. 2009. Morphology and development of the gynoecium in Centrolepidaceae: The most remarkable range of variation in Poales. *Amer. J. Bot.* 96: 1925–1940.
- Soltis, D. E. & P. S. Soltis. 2004. *Amborella* not a “basal angiosperm”? Not so fast. *Amer. J. Bot.* 91: 997–1001.
- , ———, M. E. Mort, M. W. Chase, V. Savolainen, S. B. Hoot & C. M. Morton. 1998. Inferring complex phylogenies using parsimony: An empirical approach using three large DNA sets for angiosperms. *Syst. Biol.* 47: 32–42.
- , ———, M. W. Chase, M. E. Mort, T. D. Albach, M. Zanis, V. Savolainen, W. H. Hahn, S. B. Hoot, M. F. Fay, M. Axtell, S. M. Swensen, L. M. Prince, W. J. Kress, K. C. Nixon & J. S. Farris. 2000. Angiosperm phylogeny inferred from 18S rDNA, *rbcl*, and *atpB* sequences. *Bot. J. Linn. Soc.* 133: 381–461.
- Soreng, R. J. & J. I. Davis. 1998. Phylogenetics and character evolution in the grass family (Poaceae): Simultaneous analysis of morphological and chloroplast DNA restriction site character sets. *Bot. Rev. (Lancaster)* 64: 1–85.
- Stamatakis, A. 2004. RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22: 2688–2690.
- Stevens, P. F. 2009. Angiosperm phylogeny website. <<http://www.mobot.org/MOBOT/research/APWeb/>>, accessed 5 January 2010.
- Stevenson, D. W. 1998. Mayacaceae. Pp. 294–295 in K. Kubitzki, H. Huber, P. J. Rudall, P. S. Stevens & T. Stützel (editors), *The Families and Genera of Vascular Plants*, Vol. 4. Flowering Plants, Monocotyledons: Alismatanae and Commelinanae (Except Gramineae). Springer Verlag, Berlin.
- & H. Loconte. 1995. Cladistic analysis of monocot families. Pp. 543–578 in P. J. Rudall, P. J. Cribb, D. F. Cutler & C. J. Humphries (editors), *Monocotyledons: Systematics and Evolution*. Royal Botanic Gardens, Kew, Richmond.
- Stützel, T. 1986. Die epipetalen Drüsen der Gattung *Eriocaulon* (Eriocaulaceae). *Beitr. Biol. Pflanzen* 60: 271–276.
- . 1990. “Appendices” am Gynoceum der Xyridaceen Morphogenie, Funktion und systematische Bedeutung. *Beitr. Biol. Pflanzen* 65: 275–299.
- . 1998. Eriocaulaceae. Pp. 197–207 in K. Kubitzki, H. Huber, P. J. Rudall, P. S. Stevens & T. Stützel (editors), *The Families and Genera of Vascular Plants*, Vol. 4. Flowering Plants, Monocotyledons: Alismatanae and Commelinanae (Except Gramineae). Springer Verlag, Berlin.
- Swofford, D. L. 2002. PAUP*: Phylogenetic Analysis Using Parsimony, Vers. 4.0b10. Sinauer Associates, Sunderland, Massachusetts.
- , P. J. Waddell, J. P. Huelsenbeck, P. G. Foster, P. O. Lewis & J. S. Rogers. 2001. Bias in phylogenetic estimation and its relevance to the choice between parsimony and likelihood methods. *Syst. Biol.* 50: 525–539.
- Tamura, M. N. 1998. Nartheciaceae. Pp. 381–392 in K. Kubitzki (editor), *The Families and Genera of Vascular Plants*. Springer Verlag, Berlin.
- , S. Fuse, H. Azuma & M. Hasebe. 2004. Biosystematic studies on the family Tofieldiaceae. I. Phylogeny and circumscription of the family inferred from DNA sequences of *matK* and *rbcl*. *Pl. Biol. (Stuttgart)* 6: 562–567.
- Taylor, M. & J. H. Williams. 2009. Pollination to fertilization biology of *Triphuria* (Hydatellaceae). Abstract, Botanical Society of America Summer Meeting. <<http://2009.botanyconference.org/engine/search/index.php?func=detail&aid=696>>, accessed 8 July 2010.
- Threatened Species Section. 2009. Listing statement for *Centrolepis pedderensis* (pedder bristlewort). Department of Primary Industries and Water, Tasmania. <[http://www.dpiw.tas.gov.au/inter.nsf/Attachments/SSKA-72Y8GD/\\$FILE/Centrolepis%20pedderensis.pdf](http://www.dpiw.tas.gov.au/inter.nsf/Attachments/SSKA-72Y8GD/$FILE/Centrolepis%20pedderensis.pdf)>, accessed 9 July 2010.
- Tomlinson, P. B. 1986. *The Botany of Mangroves*. Cambridge University Press, Cambridge.
- Venturelli, M. & F. Bouman. 1988. Development of ovule and seed in Rapateaceae. *Bot. J. Linn. Soc.* 97: 267–294.
- Verboom, G. A., H. P. Linder & W. D. Stock. 2004. Testing the adaptive nature of radiation: Growth form and life history divergence in the grass genus *Ehrharta* (Poaceae: Ehrhartoideae). *Amer. J. Bot.* 91: 1364–1370.
- Wang, H., M. J. Moore, P. S. Soltis, C. D. Bell, S. F. Brockington, R. Alexandre, C. C. Davis, M. Latvis, S. R. Manchester & D. E. Soltis. 2009. Rosid radiation and the rapid rise of angiosperm-dominated forests. *Proc. Natl. Acad. Sci. U.S.A.* 106: 3853–3858.
- Wedin, D. A. 1995. Species, nitrogen and grassland dynamics: The constraints of stuff. Pp. 253–262 in C. Jones & J. H. Lawton (editors), *Linking Species and Ecosystems*. Chapman and Hall, New York.
- Weller, S. G., A. K. Sakaï, A. E. Rankin, A. Golonka, B. Kutcher & K. E. Ashby. 1998. Dioecy and the evolution of pollination systems in *Schiedea* and *Alsinidendron* (Caryophyllaceae: Alsinoideae) in the Hawaiian Islands. *Amer. J. Bot.* 85: 1377–1388.

- Whitehead, D. R. 1983. Wind pollination: Some ecological and evolutionary perspectives. Pp. 97–108 in L. Real (editor), *Pollination Biology*. Academic Press, Orlando.
- Whitfield, J. B. & P. J. Lockhart. 2007. Deciphering ancient radiations. *Trends Ecol. Evol.* 22: 258–265.
- Wikström, N., V. Savolainen & M. W. Chase. 2001. Evolution of the angiosperms: Calibrating the family tree. *Proc. Roy. Soc. London, Ser. B, Biol. Sci.* 268: 2211–2220.
- Wilson, K. L. & D. A. Morrison (editors). 2000. *Monocots—Systematics and Evolution*. Proceedings of the Second International Conference on the Comparative Biology of the Monocots, Sydney. CSIRO, Melbourne.
- Wu, F.-H., D.-P. Kan, S.-B. Lee, H. Daniell, Y.-W. Lee, C.-C. Lin, N.-S. Lin & C.-S. Lin. 2009. Complete nucleotide sequence of *Dendrocalamus latiflorus* and *Bambusa oldhamii* chloroplast genomes. *Tree Physiol.* 29: 847–856.
- Wyman, S. K., R. K. Jansen & J. L. Boore. 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* 20: 3252–3255.
- Yang, Z. H. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites. *J. Molec. Evol.* 39: 306–314.
- Zomlefer, W. B. 1999. Advances in angiosperm systematics: Examples from the Liliales and Asparagales. *J. Torrey Bot. Soc.* 126: 58–62.

APPENDIX 1. Taxa included in study, with GenBank accession numbers and voucher data.

Major clade	Order	Family	Species	GenBank accession numbers*	Voucher data†
Basal angiosperms	Amborellales	Amborellaceae	<i>Amborella trichopoda</i> Baill.	NC_005086	Goremykin et al., 2003a
	Nymphaeales	Nymphaeaceae	<i>Nuphar advena</i> Aiton	NC_008788	Raubeson et al., 2007
	Austrobaileyales	Schisandraceae	<i>Illicium oligandrum</i> Merr. & Chun	NC_009600	Hansen et al., 2007
Magnoliids	Canellales	Winteraceae	<i>Drimys granadensis</i> L. f.	NC_008456	Cai et al., 2006
	Laurales	Calycanthaceae	<i>Calycanthus floridus</i> L.	NC_004993	Goremykin et al., 2003b
	Magnoliales	Magnoliaceae	<i>Liriodendron tulipifera</i> L.	NC_008326	Cai et al., 2006
Eudicots	Piperales	Piperaceae	<i>Piper cenocladum</i> Diels	NC_008457	Cai et al., 2006
	Ranunculales	Berberidaceae	<i>Nandina domestica</i> Thunb.	NC_008336	Raubeson et al., 2007
	Proteales	Platanaceae	<i>Platanus occidentalis</i> L.	NC_008335	Moore et al., 2006
	Buxales	Buxaceae	<i>Buxus microphylla</i> Siebold & Zucc.	NC_009599	Hansen et al., 2007
	Cucurbitales	Cucurbitaceae	<i>Cucumis sativus</i> L.	NC_007144	Plader et al., 2007
	Fabales	Fabaceae	<i>Medicago truncatula</i> Gaertn.	NC_003119	Matsushima et al., 2008
	Malpighiales	Salicaceae	<i>Populus alba</i> L.	NC_008235	Okumura et al., 2007
	Vitales	Vitaceae	<i>Vitis vinifera</i> L.	NC_007957	Jansen et al., 2006
	Caryophyllales	Amaranthaceae	<i>Spinacia oleracea</i> L.	NC_002202	Schmitz-Linneweber et al., 2001
	Monocots	Gentianales	Rubiaceae	<i>Coffea arabica</i> L.	NC_008535
Asterales		Asteraceae	<i>Helianthus annuus</i> L.	NC_007977	Jansen et al., 2007
Apiales		Apiaceae	<i>Anethum graveolens</i> L.	EU016721–EU016801	Jansen et al., 2007
Apiales		Araliaceae	<i>Panax ginseng</i> C. A. Mey.	NC_006290	Kim & Lee, 2004
Acorales		Acoraceae	<i>Acorus americanus</i> (Raf.) Raf.	DQ069337–DQ069702, EU0167701–EU016720	Leebens-Mack et al., 2005
Alismatales		Araceae	<i>Acorus calamus</i> L. <i>Lemna minor</i> L.	NC_007407 NC_010109	Goremykin et al., 2005 Mardanov et al., 2008
Dioscoreales		Dioscoreaceae	<i>Dioscorea elephantipes</i> (L'Hér.) Engl.	NC_009601	Hansen et al., 2007
Pandanales		Pandanaceae	<i>Pandanus utilis</i> Bory	this study*	Zomlefer 2348 (GA)
Liliales		Liliaceae	<i>Lilium superbum</i> L.	this study*	Givnish UW-8-2009-1 (WIS)
Asparagales		Amarylidaceae	<i>Agapanthus praecox</i> Willd.	this study*	Zomlefer 2311 (GA)
	Asparagaceae	<i>Albuca kirkii</i> (Baker) Brenan <i>Asparagus officinalis</i> L.	this study* this study*	McKain 111 (GA) Leebens-Mack 1001-2010 (GA)	

APPENDIX 1. Continued.

Major clade	Order	Family	Species	GenBank accession numbers*	Voucher data†
			<i>Chlorophytum rhizopendulum</i> Bjorå & Hemp	this study*	McKain 110 (GA)
			<i>Hesperaloe parviflora</i> (Torr.) J. M. Coult.	this study*	McKain 102 (GA)
			<i>Hosta ventricosa</i> (Salisb.) Stearn	this study*	McKain 106 (GA)
			<i>Lomandra longifolia</i> Labill.	this study*	Steele 1087 (UMO)
			<i>Nolina atopocarpa</i> Bartlett	this study*	McKain 114 (GA)
			<i>Yucca schidigera</i> Ortgies	DQ069337–DQ069702, EU016681–EU016700	Leebens-Mack et al., 2005
		Asteliaceae	<i>Neostelia spectabilis</i> J. B. Williams	this study*	Bruhl 2767, Quinn 95289 (NE)
		Hypoxidaceae	<i>Curculigo capitulata</i> (Lour.) Kuntze	this study*	Steele 1081 (UMO)
		Iridaceae	<i>Iris virginica</i> L.	this study*	Pires 2009-101 (UMO)
		Orchidaceae	<i>Apostasia wallichii</i> R. Br.	this study*	Zich 634; CNS130807 (CNS)
			<i>Phalaenopsis aphrodite</i> Rehb. f.	NC_007499	Chang et al., 2006
		Xanthorrhoeaceae	<i>Phormium tenax</i> J. R. Forst. & G. Forst.	this study*	Givnish Tas-2009-5 (WIS)
	Arecales	Arecaceae	<i>Chamaedorea seifrizii</i> Burret	this study*	Zomlefer 2358 (GA)
			<i>Elaeis oleifera</i> (Kunth) Cortés	EU016883–EU016962	Leebens-Mack et al., 2005
			<i>Ravenea hildebrandtii</i> C. D. Bouché	this study*	Zomlefer 2357 (GA)
	Dasygogonales	Dasygogonaceae	<i>Dasygogon bromeliifolius</i> R. Br.	this study*	KRT3702 (PERTH)
			<i>Kingia australis</i> R. Br.	this study*	KRT3703 (PERTH)
	Commelinales	Commelinaceae	<i>Belosynapsis ciliata</i> (Blume) R. S. Rao	this study*	Winters, Higgins & Higgins 186, pl 2; SI 1982-232 (SI)
			<i>Tradescantia ohiensis</i> Raf.	this study*	Moore 337 (FLA)
	Zingiberales	Musaceae	<i>Musa acuminata</i> Colla	EU016983–EU017063	Leebens-Mack et al., 2005
		Zingiberaceae	<i>Renalmia alpinia</i> (Rottb.) Maas	this study*	Zomlefer 2322 (GA)
	Poales	Bromeliaceae	<i>Brocchinia micrantha</i> (Baker) Mez	this study*	Givnish UW-8-2009-2 (WIS)
			<i>Fosterella caulescens</i> Rauh	this study*	Rauh 40573A (SEL)
			<i>Navia saxicola</i> L. B. Sm.	this study*	Givnish 3/16/1987 (WIS)
			<i>Neoregelia carolinae</i> (Beer) L. B. Sm. 'Argentea'	this study*	McKain 112 (GA)
			<i>Pitcairnia felicianae</i> (A. Chev.) Harms & Mildbr.	this study*	Luther 28 Aug 2000 (SEL)
			<i>Puya laxa</i> L. B. Sm.	this study*	Leebens-Mack 1003-2010 (GA)

APPENDIX 1. Continued.

Major clade	Order	Family	Species	GenBank accession numbers*	Voucher data†
		Centrolepidaceae	<i>Centrolepis monogyna</i> Benth.	this study*	<i>McKain 116</i> (GA)
		Cyperaceae	<i>Cyperus alternifolius</i> L.	this study*	<i>Lebens-Mack 1002-2010</i> (GA)
		Eceidiocoleaceae	<i>Eceidiocolea monostachya</i> F. Muell.	this study*	KRT3786 (PERTH)
			<i>Georgeantha hexandra</i> B. G. Briggs & L. A. S. Johnson	this study*	KRT3775 (PERTH)
		Eriocaulaceae	<i>Syngonanthus chrysanthus</i> Ruhland	this study*	<i>M. Ames 10/15/2009</i> (WIS)
		Flagellariaceae	<i>Flagellaria indica</i> L.	this study*	<i>K. Hansen 77-394</i> (BH)
		Joinvilleaceae	<i>Joinvillea ascendens</i> Gaudich. ex Brongn. & Gris	this study*	<i>Lorence 9066</i> (NTBG), <i>800379</i> (NTBG)
			<i>Joinvillea plicata</i> (Hook. f.) Newell & B. C. Stone	FJ486219–FJ486269	Leseberg & Duvall, 2009
		Juncaceae	<i>Juncus effusus</i> L.	this study*	<i>McKain 113</i> (GA)
		Mayacaceae	<i>Mayaca fluviatilis</i> Aubl.	this study*	<i>McKain 118</i> (GA)
		Poaceae	<i>Agrostis stolonifera</i> L.	NC_008591	Saski et al., 2007
			<i>Anomochloa marantoidea</i> Brongn.	NC_014062	Morris & Duvall, 2010
			<i>Bambusa oldhamii</i> Munro	NC_012927	Wu et al., 2009
			<i>Eleusine coracana</i> (L.) Gaertn.	this study*	<i>Lebens-Mack 1003-2010</i> (GA)
			<i>Hordeum vulgare</i> L.	NC_008590	Saski et al., 2007
			<i>Oryza sativa</i> L.	NC_001320	Hiratsuka et al., 1989
			<i>Puelia olyrififormis</i> (Franch.) Clayton		<i>Clayton 1060</i> (MO)
			<i>Saccharum officinarum</i> L.	NC_006084	Asano et al., 2004
			<i>Sorghum bicolor</i> (L.) Moench	NC_008602	Saski et al., 2007
			<i>Streptochaeta angustifolia</i> Soderstr.	this study*	<i>J. I. Davis 757</i> (BH)
			<i>Triticum aestivum</i> L.	NC_002762	Ogihara et al., 2002
			<i>Zea mays</i> L.	NC_001666	Maier et al., 1995
		Rapateaceae	<i>Potarophytum riparium</i> Sandwith	this study*	<i>Givnish GUY-09-2</i> (BRG)
		Restionaceae	<i>Thamnochortus insignis</i> Mast.	this study*	<i>Givnish UW-8-2009-3</i> (WIS)
		Sparganiaceae	<i>Sparganium eurycarpum</i> Engelm.	this study*	<i>Ames 10/21/2009</i> (WIS)
		Thurniaceae	<i>Thurnia sphaerocephala</i> Hook. f.	this study*	<i>Givnish GUY-09-5</i> (BRG)
		Typhaceae	<i>Typha latifolia</i> L.	NC_013823	Guisinger et al., 2010
		Xyridaceae	<i>Abolboda macrostachya</i> Spruce ex Malme	this study*	<i>Givnish GUY-09-7</i> (BRG)

* GenBank accession numbers for plastid genes newly sequenced in this study are HQ180399–HQ183709. A spreadsheet listing individual accession number for each region and species is available at <<http://chloroplast.cbio.psu.edu/supplement.html>>.

† Voucher specimen (collector and number, with acronym for herbarium of deposit, or citation for sequences previously published elsewhere).

APPENDIX 2. Habitat and floral characteristics of taxa included in tests of correlated evolution involving anemophily.

ECOLOGICAL CHARACTERISTICS

Data on the ecology of each lineage (Table A1) were drawn from summaries in Dahlgren et al. (1985), Givnish et al. (1999, 2000, 2004, 2007), Linder and Rudall (2005), and Sokoloff et al. (2009), as well as a variety of other sources noted below. For small, ecologically uniform lineages (e.g., Centrolepidaceae, Eriocaulaceae, Xyridaceae), such scoring was straightforward. Where needed in large, ecologically diverse families or orders represented in our phylogenetic analysis by single taxa, we overlaid the ancestral conditions previously inferred for such groups based on detailed molecular studies (e.g., Givnish et al., 2000, 2004, 2007), or inferred what those ancestral conditions would have been, given the habitats of the earliest divergent lineages within those families or orders.

Given their distinctive ecologies, we separated subfamily Anomochloideae from the other Poaceae sampled (a subset of the bistigmatic clade = subfamily Puelioideae + the PACCMAD-BEP clade of Sánchez-Ken et al., 2007). We interpolated the subfamily Pharioideae (unsampled in our phylogenetic study) between Anomochloideae and Puelioideae, given that recent molecular studies place them there with high support, and given the distinctive ecology of pharoids and puelioids versus most members of the huge PACCMAD-BEP clade, which includes 99% of all grass genera and species (Soderstrom & Calderón, 1971; Clark et al., 1995, 2000; Grass Phylogeny Working Group I, 2001; Hodkinson et al., 2007a, b; Bouchenak-Khelladi et al., 2008). The first three subfamilies all occur on shaded, well-drained, fire-protected, nonsandy soils in tropical rainforest understories. With the exception of the bambusoids, many members of the PACCMAD-BEP clade grow in open, well-drained sites frequently swept by fire. Based on the multilocus phylogeny of Bouchenak-Khelladi et al. (2008)—the most comprehensive and most strongly supported tree for grasses to date, but still missing representatives of two small subfamilies (Aristoideae and Micrairoideae)—open habitats appear to be ancestral to this clade, given that such habitats characterize almost all chloridoideae, danthonioids, arundinoideae, and panicoids, as well as *Thysanolaena* Nees—*Cenotheca* Desv. sister to the panicoids in the PACCMAD clade, and almost all ehrhartoideae and pooids in the BEP clade. Bamboos and taxa scattered throughout the other subfamilies occur primarily in shaded habitats. Typical soils appear to be ancestral for the PACCMAD-BEP clade, given that extremely infertile, sandy soils characterize only *Ehrharta* Thunb. among the early divergent taxa (Verboom et al., 2004). Similarly, well-drained soils and fireswept conditions appear to be ancestral. Among early divergent members of the BEP clade, only *Oryza* L. and *Glyceria* R. Br. are wetland lineages; among early divergent members of the PACCMAD clade, the initial arundinoid and danthonioid groups successively sister to the chloridoideae that grow in wetlands, while the panicoids and centothecoids generally grow in uplands. Parsimony thus implies that well-drained conditions are ancestral to the PACCMAD-BEP clade as a whole. Only *Oryza*, bamboos, and *Merxmüllera* Conert appear not to be favored by fire among the early divergent lineages, implying that fireswept conditions are also ancestral.

Joinvillea and *Flagellaria* occur on shaded, well-drained, reasonably fertile sites in tropical rainforests and generally do not experience fire (Backer, 1951; Newell, 1969; Linder, 1987; Bayer & Appel, 1998; Linder & Rudall, 2005).

Eceidocoleaceae are native to open, well-drained, fireswept sites on extremely infertile soils in southwest Australia (Linder et al., 1998; Linder & Rudall, 2005). Most Eriocaulaceae and Xyridaceae inhabit similar but frequently wet or inundated sites worldwide (Kral, 1998; Stützel, 1998; Linder & Rudall, 2005). Restionaceae occur on open, extremely infertile, and frequently fireswept sites in the Southern Hemisphere, especially in southwestern Africa and southwest Australia; different groups inhabit well-drained and inundated microsites (Linder & Rudall, 2005). Based on the molecular phylogeny of Linder et al. (2003), wet to inundated sites appear to be ancestral, given their occupancy by one Australian clade (consisting of *Sporodanthus* F. Muell. ex J. Buch., *Calorophus* Labill., and *Lepyrodia* R. Br.), and by early divergent members (i.e., *Winifredia* L. A. S. Johnson & B. G. Briggs, *Empodisma* L. A. S. Johnson & B. G. Briggs) of a second Australian clade (see Jordan, 2009). The molecular phylogeny of Briggs et al. (2000) also implies that wet to inundated microsites were ancestral for Restionaceae, given their occupancy by the Australian *Sporodanthus*–*Calorophus*–*Lepyrodia* clade (now Sporodanthoideae) and by early divergent members (i.e., *Eurychorda* B. G. Briggs & L. A. S. Johnson, *Empodisma*, *Winifredia*, *Taraxis* B. G. Briggs & L. A. S. Johnson) of the second Australian clade (now Leptocarpoideae). Centrolepidaceae grow on open, extremely infertile, sparsely vegetated sites, in ephemerally wet or permanently sodden microsites, ranging from near sea level to wet alpine turfs (Pignatti & Pignatti, 1994, 2005; Cooke, 1998, 2010; Threatened Species Section, 2009). Such sites are very unlikely to be flammable (Linder & Rudall, 2005). *Mayaca* grows in open, seasonally, and permanently inundated sites on extremely infertile substrates (Stevenson, 1998) and is generally immune to fire (Linder & Rudall, 2005).

We used the phylogenies produced by Simpson et al. (2003), Drábková and Vlček (2009), and Muasya et al. (2009) to infer that sunny habitats are ancestral to all families of the cyperids Thurniaceae–(Cyperaceae–Juncaceae), based on the occupancy of such sites (1) by both monotypic genera of Thurniaceae; (2) by *Juncus trifidus* L., *J. monanthos* Jacq., *Luzula subcongesta* (S. Watson) Jeps., *L. piperi* (Coville) M. E. Jones, *J. potaninii* Buchenau, *J. oxycarpus* E. Mey. ex Kunth, *J. stygius* L., and other taxa sister to all remaining members of Juncaceae, or of the individual early divergent clades of Juncaceae; and (3) by the *Coleochloa* Gilly–*Microdracooides* Hua–*Trilepis* Nees complex, *Cladium* P. Browne, and *Gymnoschoenus* Nees, which are sister successively to all other members of Cyperaceae subfamily Cyperioideae, in contrast to members of sister subfamily Mapanioideae, which mostly grow in forest understories. Note that none of the taxa just mentioned (except *Thurnia* of Thurniaceae) were included in our plastome sequencing study. Given the basal split between Cyperioideae and Mapanioideae, we coded Cyperaceae as ambiguous (i.e., polymorphic) for sunny/shady habitats; Juncaceae and Thurniaceae were coded as sunny. Thurniaceae occur on extremely infertile, seasonally inundated substrates in the Guayana Shield (*Thurnia*) and South Africa (*Prionium*). These sites are clearly subject to fire in *Prionium* (charred petiole bases and stems, T. J. Givnish, pers. obs.) but probably are not in the largely aquatic *Thurnia*. Cyperaceae was scored as polymorphic for soil fertility and fire frequency, given that the earliest divergent clades in Trilepideae occur on highly infertile substrates (granitic inselbergs, Everglades and similar habitats, button-grass moorlands), the first two of which are fireswept, while more fertile soils and a general absence of fire characterize Mapanioideae. Well-drained substrates were scored as

Table A1. Habitat characteristics assigned to Poalean ingroups and outgroups for character-state reconstruction. See appendix text for data sources.

	Light availability*	Moisture supply†	Nutrient supply‡	Fire prevalence§
Poaceae, PACCMAD-BEP clade	1	0	1	1
Poaceae, Puelioideae	0	0	1	0
Poaceae, Pharioideae	0	0	1	0
Poaceae, Anomochlooideae	0	0	1	0
Ecdeiocoleaceae	1	0	0	0
Joinvilleaceae	0	0	1	0
Flagellariaceae	0	0	1	0
Restionaceae	1	0/1	0	1
Centrolepidaceae	1	0/1	0	0
Xyridaceae	1	1	0	1
Mayacaceae	1	1	0	0
Eriocaulaceae	1	1	0	1
Cyperaceae	0/1	0/1	1	0/1
Juncaceae	1	1	1	1
Thurniaceae	1	1	0	0
Rapateaceae	1	1	0	1
Typhaceae	1	1	1	1
Bromeliaceae	1	1	0	1
Commelinales	1	1	0	0
Zingiberales	0	0	1	0
Dasygogonales	1	0	0	1
Arecales	0/1	0/1	1	0/1
Orchidaceae	0	0	0	0

* 0 = Shaded habitats; 1 = sunny habitats.

† 0 = Well-drained microsites; 1 = seasonally or permanently inundated.

‡ 0 = Nutrient-poor sands or peats; 1 = fertile substrates.

§ 0 = Fires rare/absent; 1 = fires frequent relative to rate of plant regrowth.

ancestral, given their occurrence in mapanioids and the *Coleochloa–Microdracoides–Trilepis* clade. Well-drained substrates and lack of fire are also ancestral in Juncaceae, given their occurrence in *J. monanthos* and *J. trifidus* L. and in *Luzula* (the two earliest divergent rush lineages, fide Drábková & Vlček, 2009). Substrate fertility was scored as polymorphic, given extremely infertile substrates (i.e., near lack of soil) in *J. monanthos* and *J. trifidus* and richer soils in *Luzula* and *Juncus* L. subg. *Juncus* (Table A1).

All habitats inhabited by Rapateaceae are exceedingly infertile, and almost all are open (except for those occupied by *Epidryos* Maguire, *Monotrema* Körn., *Rapatea* Aubl., and *Windsorina* Gleason), wet to inundated (save those occupied by *Epidryos* and *Rapatea*), and fireswept (but for those occupied by *Epidryos*, *Monotrema*, *Rapatea*, *Windsorina*, and possibly the unstudied *Maschalocephalus* Gilg & K. Schum.) (Givnish et al., 2000, 2004). Based on the phylogeny presented by Givnish et al. (2004), open, wet, fireswept, and highly infertile sites are thus ancestral for Rapateaceae. *Typha* L. and *Sparganium* L. of Typhaceae both occur on open, wet to inundated, fertile substrates that rarely if ever burn (Linder & Rudall, 2005). Ancestral habitats for Bromeliaceae appear to have been open, wet, and exceedingly infertile, based on the restriction to such sites of most species of *Brocchinia* and *Lindmania* and the current multilocus phylogeny for that family (see Givnish et al., 1997, 2007). We scored bromeliads as being polymorphic for fire frequency (Table A1), given that several early divergent members of *Brocchinia* (e.g., *B. prismatica*, *B. melanacra*) occur in fireswept habitats, while members of *Lindmania*

occur in both fireswept habitats (e.g., *L. guianensis*) and those that appear to burn rarely if at all (e.g., *L. arachnoidea* (L. B. Sm., Steyerl. & H. Rob.) L. B. Sm.); members of the third divergent subfamily Tillandsioideae (Givnish et al., in prep.) rarely experience fire.

Among the outgroups, ancestral habitats for Commelinales were scored as open, wet, nutrient poor, and fire immune, based on the sharing of these characteristics by *Hanguana* Blume, *Cartonema* R. Br., *Philydrella* Caruel, and *Anigozanthos* Labill./*Xiphidium* Aubl. and on the phylogeny of Saarela et al. (2008). By comparison, the ancestral habitats of its sister group Zingiberales were scored as closed, well drained, fertile, and fire immune, based on these environments being shared by present-day Lowiaceae, *Heliconia* L./*Musa* L. and the “ginger” families (Cannaceae, Marantaceae, Costaceae, Zingiberaceae), and on the relationships of these families to each other and Strelitziaceae in the multilocus phylogeny of Johansen (2005). Present-day habitats of Dasygogonaceae are uniformly open, well drained, exceedingly infertile, and fireswept (Clifford et al., 1998). Ancestral habitats of Arecales were scored as shaded, well drained, and neither extremely infertile nor fireswept, based on the supermatrix and supertree phylogenies presented by Baker et al. (2009) and the sharing of such habitats today by most calamoids and coryphoids. Exceptions to these rules are numerous; for example, in the predominantly rainforest-dwelling Calamoideae (the rattans and their allies), *Mauritia* L. f. and related genera occur in open, seasonally inundated, often fireswept palm savannas. Finally, ancestral habitats for Orchidaceae were scored as shaded, well drained, moder-

Table A2. Habitat and floral characteristics of taxa included in tests of correlated evolution. See text for data sources.

	Wind pollina- tion	Habitat open	Dioecy	Nectar	Ovules multiple	Flowers unisexual	Flowers large	Flowers showy
<i>Abolboda</i> Bonpl.	0	1	0	1	1	0	1	1
<i>Agrostis</i> L.	1	1	0	0	0	1	0	0
<i>Anomochloa</i> Brongn.	0	0	0	0	0	0	0	0
<i>Apostasia</i> Blume	0	0	0	0	1	0	1	1
<i>Bambusa</i> Schreb.	1	0	0	0	1	1	0	0
<i>Belosynapsis</i> Hassk.	0	0	0	0	1	0	1	1
<i>Brocchinia</i> Schult. f.	0	1	0	1	1	0	1	1
<i>Centrolepis</i> Labill.	1	1	0	0	0	1*	0	0
<i>Chamaedorea</i> Willd.	0	0	1	1	0	1	0	0
<i>Curculigo</i> Gaertn.	0	0	0	0	1	0	1	1
<i>Cyperus</i> L.	1	1	0	0	0	1	0	0
<i>Dasyogon</i> R. Br.	0	1	0	1	0	0	1	1
<i>Ecdiocolea</i> F. Muell.	1	1	0	0	0	1	0	0
<i>Elaeis</i> Jacq.	1	1	0	0	0	1	0	0
<i>Eleusine</i> Gaertn.	1	1	0	0	0	1	0	0
<i>Flagellaria</i> L.	0	0	0	1	0	0	0	1
<i>Fosterella</i> L. B. Sm.	0	0	0	1	1	0	0	1
<i>Georgeantha</i> B. G. Briggs & L. A. S. Johnson	1	1	0	0	0	1	0	0
<i>Hordeum</i> L.	1	1	0	0	0	1	0	0
<i>Joinvillea-1</i> Gaudich. ex Brongn. & Gris	0	0	0	0	0	0	0	0
<i>Joinvillea-2</i>	0	0	0	0	0	0	0	0
<i>Juncus</i> L.	1	1	0	0	1	0	0	0
<i>Kingia</i> R. Br.	0	1	0	1	1	0	1	1
<i>Mayaca</i> Aubl.	0	1	0	0	1	0	1	1
<i>Musa</i> L.	0	0	0	1	1	0	1	1
<i>Navia</i> Schult. f.	0	1	0	1	1	0	1	1
<i>Neostelia</i> J. B. Williams	0	1	0	1	1	0	1	1
<i>Neoregelia</i> L. B. Sm.	0	0	0	1	1	0	1	1
<i>Oryz</i> L.	1	1	0	0	0	0	0	0
<i>Phalaenopsis</i> Blume	0	0	0	1	1	0	1	1
<i>Pitcairnia</i> L'Hér.	0	0	0	1	1	0	1	1
<i>Potarophytum</i> Sandwith	0	0	0	0	0	0	1	1
<i>Puelia</i> Franch.	1	0	0	0	0	0	0	0
<i>Puya</i> Molina	0	1	0	1	1	0	1	1
<i>Ravenea</i> C. D. Bouché	0	0	0	1	0	0	0	0
<i>Renealmia</i> L. f.	0	0	0	1	1	0	1	1
<i>Saccharum</i> L.	1	0	0	0	0	0	0	0
<i>Sorghum</i> Moench	1	1	0	0	0	0	0	0
<i>Sparganium</i> L.	1	1	0	0	0	1	0	0
<i>Streptochaeta</i> Schrad. ex Nees	0	0	0	0	0	0	0	0
<i>Syngonanthus</i> Ruhland	0	1	0	1	1	1	1	1
<i>Thamnochortus</i> P. J. Bergius	1	1	1	0	0	1	0	0
<i>Thurnia</i> Hook. f.	1	1	0	0	1	1	0	0
<i>Tradescantia</i> L.	0	1	0	0	1	0	1	1
<i>Triticum</i> L.	1	1	0	0	0	1	0	0
<i>Typha</i> L.	1	1	0	0	0	1	0	0
<i>Zea</i> L.	1	1	0	0	0	1	0	0

* The traditional interpretation is that centrolepid flowers are unisexual, but Sokoloff et al. (2010) argue instead that the “inflorescence” is a highly reduced hermaphroditic flower.

ately fertile, and unlikely to burn, based on the sharing of these characteristics in present-day habitats of *Apostasia/Neuwiedia* Blume of Apostasioideae, and *Selenipedium* Rehb. f. and some early divergent *Cyripedium* L. (i.e., *C.*

irapeanum La Llave & Lex., *C. plectrochilum* Franch., *C. fasciculatum* Kellogg) of Cyripedioideae, in light of molecular phylogenies for the family (Cox et al., 1997; Cameron, 2004; Górniak et al., 2010).

CORRELATED EVOLUTION BETWEEN WIND POLLINATION AND
VARIOUS PLANT TRAITS

We conducted tests for patterns of correlated evolution of wind pollination in relation to plant habitat, nectar secretion, floral sexuality, sexual system, ovule number, floral size, and floral showiness (see text for scoring, analyses, and references). Characters were scored for the taxa whose plastomes were sequenced (Table A2). Long-branched or plumose stigmas (sensu Heslop-Harrison & Shivanna, 1977; Dahlgren et al., 1985; Soreng & Davis, 1998) were taken as one line of evidence for wind pollination and consequently not scored separately as a character that might show correlated evolution with wind pollination. Linder (1987) and Linder and Rudall (2005) considered *Flagellaria* and *Joinvillea* to be wind pollinated, but we reject that view. Both genera lack plumose stigmas (Soreng & Davis, 1998). *Flagellaria* has fragrant white flowers (Backer, 1951), and ants visit *Joinvillea* flowers (Newell, 1969) while the plant secretes large amounts of nectar through extrafloral nectaries (Blüthgen & Fiedler, 2004; Blüthgen et al., 2004a, b). *Anomochloa*, *Streptochoeta*, and *Pharus*—as well as several herbaceous bamboos that are also native to rainforest understories—all appear to be animal pollinated, contrary to Linder and Rudall (2005), based on direct observations in some cases and on these taxa possessing simple stigmas and pollen with sculpture divergent from that typical of wind-pollinated grasses (Soderstrom & Calderón, 1971, 1979; Soderstrom, 1981; Soreng & Davis, 1998). As a conservative

measure, *Puelia* Franch. was scored as wind pollinated, reflecting its possession of plumose stigmas (Grass Phylogeny Working Group I, 2001). Nothing is known about the pollination biology of *Puelia* and its sister *Guaduella* Franch., however, and their habit and growth in windless forest understories are strikingly similar to those of *Anomochloa*, *Streptochoeta*, and *Pharus*. Possession of plumose stigmas is not, however, an unfailing indicator of wind pollination—the herbaceous Neotropical bamboo *Pariana* is insect pollinated (Soderstrom & Calderón, 1971) but also has plumose stigmas. *Guaduella* has white, visually conspicuous anthers. If *Puelia* and *Guaduella* were instead insect pollinated, the significance of several of our tests for correlated evolution involving wind pollination would increase.

The extension of the scoring of wind versus biotic pollination from individual genera (Table A2) to families/orders is straightforward in all cases except Eriocaulaceae and Arecales. For Eriocaulaceae, few rigorous studies on the mechanism of pollination have been conducted; those that have (i.e., Stützel, 1986; Rosa & Scatena, 2003, 2007; Ramos et al., 2005; Oriani et al., 2009) have identified biotic pollination as being involved. Thus, even though many previous authors have scored Eriocaulaceae as being wind pollinated, we score the family as animal pollinated. For palms, it is clear that the rattans, *Mauritia*, and *Nypa* Steck are all insect pollinated (Tomlinson, 1986; Ervik, 1993; Heard, 1999). Given their position in the current molecular phylogeny for Arecaeae (Baker et al., 2009), we conclude that animal pollination is the ancestral condition for the palm family.