# Epigenetic Silencing of Host Cell Defense Genes Enhances Intracellular Survival of the Rickettsial Pathogen *Anaplasma phagocytophilum*

# Jose C. Garcia-Garcia<sup>1</sup>, Nicole C. Barat<sup>1</sup>, Sarah J. Trembley<sup>2</sup>, J. Stephen Dumler<sup>1\*</sup>

1 Department of Pathology, The Johns Hopkins University School of Medicine, Baltimore, Maryland, United States of America, 2 Department of Molecular and Comparative Pathobiology, The Johns Hopkins University School of Medicine, Baltimore, Maryland, United States of America

## Abstract

Intracellular bacteria have evolved mechanisms that promote survival within hostile host environments, often resulting in functional dysregulation and disease. Using the *Anaplasma phagocytophilum*–infected granulocyte model, we establish a link between host chromatin modifications, defense gene transcription and intracellular bacterial infection. Infection of THP-1 cells with *A. phagocytophilum* led to silencing of host defense gene expression. Histone deacetylase 1 (HDAC1) expression, activity and binding to the defense gene promoters significantly increased during infection, which resulted in decreased histone H3 acetylation in infected cells. HDAC1 overexpression enhanced infection, whereas pharmacologic and siRNA HDAC1 inhibition significantly decreased bacterial load. HDAC2 does not seem to be involved, since *HDAC2* silencing by siRNA had no effect on *A. phagocytophilum* intracellular propagation. These data indicate that HDAC up-regulation and epigenetic silencing of host cell defense genes is required for *A. phagocytophilum* infection. Bacterial epigenetic regulation of host cell gene transcription could be a general mechanism that enhances intracellular pathogen survival while altering cell function and promoting disease.

Citation: Garcia-Garcia JC, Barat NC, Trembley SJ, Dumler JS (2009) Epigenetic Silencing of Host Cell Defense Genes Enhances Intracellular Survival of the Rickettsial Pathogen Anaplasma phagocytophilum. PLoS Pathog 5(6): e1000488. doi:10.1371/journal.ppat.1000488

Editor: Craig R. Roy, Yale University School of Medicine, United States of America

Received December 30, 2008; Accepted May 27, 2009; Published June 19, 2009

**Copyright:** © 2009 Garcia-Garcia et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This research was supported by NIH research grant R01-Al44102 to JSD from the National Institutes of Allergy and Infectious Diseases. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

\* E-mail: sdumler@jhmi.edu

## Introduction

Intracellular pathogens, through a long-standing association with host cells, have evolved mechanisms that allow survival within the often hostile environment of their hosts [1]. These mechanisms usually result in dramatic transcriptional changes in infected host cells and in dysregulation of cell functions that potentially lead to disease. Global analysis of mammalian gene expression in response to intracellular bacteria has led to the identification of major pathways affected during infection [2]. Due to the limited genetic and metabolic resources of intracellular bacteria, these pathogens likely evolved global and efficient mechanisms for host cell gene regulation. While signaling pathways and transcriptional regulators often act on a limited subset of genes, epigenetic regulators tend to more globally control gene expression, and impact major cellular processes such as cell cycle progression and cell differentiation. Dysregulation of epigenetic control mechanisms often leads to dramatic phenotypic changes.

Reversible histone acetylation is a key epigenetic regulator of chromatin structure and gene expression, in combination with other posttranslational modifications. These patterns of histone modification are maintained by histone modifying enzymes such as histone deacetylases (HDAC). Disruption of HDAC activity with inhibitors or by siRNA affects expression of up to 10% of the genes in different cell types [3–6]. Global HDAC-mediated transcriptional changes can have a concomitant effect on cell function – an epigenetic mechanism often exploited by viruses to promote infection [7–9]. Recent reports show that intracellular bacteria manipulate host cell epigenetics to facilitate infection as well [10-12].

The tick-transmitted rickettsial pathogen Anaplasma phagocytophilum, causative agent of human granulocytic anaplasmosis, is one of only four bacteria known to survive and propagate within human neutrophils and their bone marrow progenitors. Neutrophils are generally considered unsuitable host cells for intracellular bacteria because they are short-lived and are primary defense cells equipped with diverse antimicrobial mechanisms. Major neutrophil functions are altered with A. phagocytophilum infection, which ultimately results in clinical disease. Processes such as oxidative burst, apoptosis and phagocytosis are inhibited or delayed by A. phagocytophilum infection [13-15], while degranulation and cytokine/chemokine production are activated [16-18]. These dramatic alterations in host cell function can be explained at least in part by A. phagocytophilum-induced host transcriptional changes [19-22]. We have recently shown that the A. phagocytophilum effector protein AnkA is translocated into the host nucleus, where it interacts with host chromatin to affect gene transcription [23-25]. However, global mechanisms leading to A. phagocytophilum-induced transcriptional changes remain poorly defined.

In this report, we investigate the hypothesis that global downregulation of key host defense genes is critical for *A. phagocytophilum* 

## **Author Summary**

Although the main function of defense cells is to eliminate invading infections, some intracellular bacterial pathogens manage to turn defense cells into suitable hosts for bacterial propagation. In doing so, intracellular pathogens dysregulate host cell function and cause disease. With genomic and metabolic resources thousands of times more limited than the host's, intracellular bacteria have evolved very efficient mechanisms to globally subvert the host defense. Here, we define a mechanism by which the intracellular pathogen Anaplasma phagocytophilum globally inhibits host cell defenses by affecting mechanisms of epigenetic control of defense gene expression. Silencing or inhibition of the host protein HDAC1 has a negative effect on intracellular bacterial replication, whereas HDAC1 overexpression leads to defense gene silencing and facilitates intracellular bacterial survival. This study not only provides new insight into a mechanism of host cell subversion, but also identifies a potential target for future development of novel therapeutic intervention strategies.

intracellular infection and propagation. We establish a link between intracellular bacterial survival/growth and changes in host transcription and function known to be involved in A. *phagocytophilum* pathogenesis. These data suggest a global epigenetic mechanism by which bacterial pathogens interact with and control host cells.

# Results

# Defense gene expression is down-regulated in *A. phagocytophilum*-infected cells

Previous studies suggest that A. phagocytophilum infection downregulates the expression of key host defense genes such as CYBB, RAC2, MPO and BPI [26]. Expression of 23 defense genes, including genes encoding for antimicrobial peptides as well as genes involved in enzymatic and oxidative defense mechanisms, was compared by quantitative RT-PCR (qRT-PCR). Expression of 19 of these defense genes was down-regulated during infection of THP-1 cells with A. phagocytophilum (Fig. 1), whereas expression of IL8 and FTH, two genes known to be upregulated during A. phagocytophilum infection of granulocytes, was increased. Notably, the EPX and MPO genes, which form a gene cluster in the genome (17q23), were down-regulated, as were most of the defensins, which comprise a gene cluster on chromosome 8p23, and the genes AZU1, ELA2 and PRTN3, which form a third defense gene cluster on chromosome 19p13 (Fig. 1). From this and previously published data, it can be



**Figure 1. Down-regulation of host defense genes with** *A. phagocytophilum* **infection of THP-1 cells.** RNA was extracted from infected and uninfected THP-1 cells 48 hours post-infection and expression of defense genes was quantitated by qRT-PCR. *IL8* and *FTH* expression were used as up-regulation controls. Gene expression changes were expressed as transcription fold-change in infected cells with respect to uninfected cells. Numbers <1 denote down-regulation and >1 indicate up-regulation. doi:10.1371/journal.ppat.1000488.g001



Anaplasma Host Epigenetics

**Figure 2. The pattern of histone post-translational modifications and HDAC1 binding in the defense gene promoters is affected by** *A. phagocytophilum* infection of THP-1 cells. Chromatin from infected and uninfected THP-1 cells was prepared 48 hours post-infection. The histone modification pattern and HDAC1 binding at the defense gene promoters was analyzed by ChIP using antibodies specific for (A) Ac-H3, (B) Me-H3 and (C) HDAC1. Immunoprecipitated DNA fragments were quantitated by qPCR using primers specific for each promoter region. Changes were expressed as the ratio of immunoprecipitated chromatin target from infected to uninfected cells enriched with respect to input chromatin. doi:10.1371/journal.ppat.1000488.g002

concluded that defense gene expression is globally downregulated with *A. phagocytophilum* infection, permitting sustained inhibition of antimicrobial defense and facilitating establishment of intracellular infection.

# *A. phagocytophilum* infection affects defense gene chromatin structure

Silencing of genes or gene clusters is often associated with changes in chromatin structure mediated by epigenetic regulators. In order to identify possible chromatin alterations associated with A. phagocytophilum infection, a chromatin immunoprecipitation (ChIP) approach was used to study alterations in histone H3 diacetylation (Ac-H3) and monomethylation (Me-H3) patterns at defense gene promoters (Fig. 2). A decrease in the Ac-H3 with A. phagocytophilum infection was observed in 9 out of 11 defense gene promoters (Fig. 2A), consistent with decreased transcriptional activity at these loci during infection (Fig. 1A, B). Slightly increased Ac-H3 was observed in the GNLY and DCD promoters, whereas no changes were observed in the IL8 promoter, which was used as a control. IL8, unlike defense genes, is upregulated by signal transduction pathways activated with infection. A concomitant increase in Me-H3 was observed for all genes except IL8. This difference was not significant for DEFA1, DEFA6 and DCD. These results suggest that silencing of most key host defense genes with A. phagocytophilum infection could occur by epigenetic changes in chromatin structure and histone post-translational modification patterns.

Since histone deacetylases play a crucial role in regulating histone post-translational modifications, we next investigated changes in HDAC1 binding to the promoter regions of host defense genes during *A. phagocytophilum* infection (Fig. 2C). After infection, HDAC1 binding was increased to the promoters of all 11 defense genes analyzed. HDAC1 acts as a transcriptional repressor and its increased association with defense gene promoters upon *A. phagocytophilum* infection suggests a role for HDAC1 in defense gene silencing.

# Deacetylase expression and activity are increased with *A. phagocytophilum* infection

Since histone deacetylases play a crucial role in regulating histone post-translational modifications, we next investigated changes in HDAC expression and activity associated with A. phagocytophilum infection. Gene expression analysis of HDAC1 and HDAC2 in A. phagocytophilum-infected cells by qRT-PCR showed a transient increase in HDAC1 expression that peaked within 24 hours post-infection, whereas HDAC2 transcription steadily increased over time (Fig. 3A). Following the initial increase, HDAC1 mRNA levels decreased by 72 hours, similar to the profile observed in A. phagocytophilum-infected HL-60 cells (not shown), suggesting HDAC1-mediated transcriptional autorepression. HDAC1 protein levels started to increase within 24 hours after A. phagocytophilum infection, increasing greater than 1.5-fold by 48 hours and moderately decreasing thereafter (Fig. 3B). HDAC2 protein levels steadily increased over time with kinetics similar to that of HDAC2 gene transcription (Fig. 3A, B). HDAC activity was also significantly increased in infected cells 48 h post-infection (Fig. 3C). Taken together, these results indicate that HDAC1 and HDAC2 expression is increased upon *A. phagocytophilum* infection by mechanisms regulated at least in part at the transcriptional level. Three human strains of *A. phagocytophilum* from North America and Europe showed similar abilities to increase HDAC expression and activity (Figure S1).

# Increased HDAC activity down-regulates defense gene expression and promotes *A. phagocytophilum* infection of THP-1 cells

To determine whether increased HDAC1 expression is required for *A. phagocytophilum* to establish successful intracellular infection, HDAC activity was inhibited prior to infection using the HDAC inhibitors trichostatin A (TSA) and sodium butyrate. *A. phagocytophilum* infection was reduced in a dose-dependent manner by both HDAC inhibitors (Fig. 4A, B). *A. phagocytophilum* infection was not significantly reduced at lower doses of either inhibitor, but at higher doses *A. phagocytophilum* infection was significantly reduced (p<0.05). Preincubation of *A. phagocytophilum* with TSA did not affect its ability to infect THP-1 cells (Fig. 4C), ruling out HDAC inhibitor toxicity toward *A. phagocytophilum*.

Because TSA inhibits not only HDAC1 but also other class I acetylases, HDAC1 expression was specifically targeted using siRNA. HDAC1 silencing resulted in a significant reduction in A. phagocytophilum load in infected THP-1 cells, whereas HDAC2 silencing did not affect A. phagocytophilum infection (Fig. 5A). The possible role of other deacetylases and the lower efficiency of siRNA compared to pharmacological inhibition, may explain the less dramatic effect of HDAC1 silencing on A. phagocytophilum infection as compared to HDAC inhibition with TSA or sodium butyrate. On the other hand, the infectious load of THP-1 cells transfected with an HDAC1-expressing plasmid increased (Fig. 5B), indicating that HDAC1 expression enhances A. phagocytophilum infection and propagation. To confirm that HDAC1 plays a direct role in silencing host defense genes, the expression of four defense genes was analyzed in THP-1 cells transfected with HDAC1 siRNA or pHDAC1-FLAG plasmid. Expression of DEFA1, AZU1, LYZ and MPO was upregulated by siRNA HDAC1 silencing, whereas A. phagocytophilum infection and HDAC1 overexpression led to decreased gene expression (Fig. 5C). These data confirm the role of HDAC1 in host defense gene silencing and host cell conditioning for intracellular survival

# Discussion

The intracellular environment is hostile for bacteria that propagate within defense cells such as macrophages, monocytes and granulocytes. These cells are equipped with an arsenal of antimicrobial molecules that target pathogens using diverse mechanisms, including generation of reactive oxygen intermediates and secretion of antimicrobial peptides and enzymes. With a cytoplasm and a genome over a thousand times smaller than those of the host cell, *A. phagocytophilum* and other intracellular bacteria must require efficient mechanisms for survival and persistence within the host cell by inducing sustained changes in host cell function. The data presented here point to a



**Figure 3. HDAC expression and activity is increased by** *A. phagocytophilum* **infection.** (A) RNA from infected and uninfected cells was extracted and *HDAC1* and *HDAC2* expression was quantitated by qRT-PCR. Transcription fold-change with respect to uninfected cells was calculated. (B) HDAC1 and HDAC2 in infected and uninfected THP-1 cells were detected by immunoblotting. Band intensity was determined by densitometric analysis and protein expression level changes with respect to the initial expression level in uninfected cells were calculated. Samples were normalized for  $\beta$ -actin content. The example shown is representative of 3 separate experiments with similar results. (C) HDAC activity in nuclear extracts of infected and uninfected cells at 48 hpi was determined using a fluorescent assay kit. Nuclear extract from HeLa cells was used as control. doi:10.1371/journal.ppat.1000488.g003

mechanism, conserved among human isolates of A. phagocytophilum, for control of host cell gene expression and function based on host cell epigenetic changes. Increased HDAC expression in A. phagocytophilum-infected cells results in accumulation of deacetylated histones. These changes in the pattern of histone posttranslational modifications likely have a direct effect on gene expression by affecting chromatin structure, leading to a highly compact chromatin conformation with limited access to transcriptional activators. Importantly, defense genes are often organized in chromosomal clusters that permit coordinate expression and regulation by changes in chromatin organization [27]. Moreover, increased HDAC expression could also directly down-regulate the expression of genes which are transcriptionally repressed by HDACs, such as CYBB [28]. The data presented here suggest a new mechanism mediated by HDACs by which CYBB and other defense genes can be silenced by A. *phagocytophilum* infection. Other bacteria seem to take advantage of similar epigenetic mechanisms of control of host cell function. Listeria monocytogenes, Clostridium perfringens and Streptococcus pneumonia induce H3 dephosphorylation as well as H4 deacetylation during infection, which correlate with down-regulated expression of a subset of host genes including defense genes [11]. Similarly, mycobacterial infection of human cells results in host gene silencing using a mechanism that involves HDAC complex formation and histone deacetylation [29]. Host cell epigenetics, bacterial effectors such as the AnkA protein, signaling pathways and other mechanisms of transcriptional regulation likely contribute to the overall regulation of host cell gene expression and function.

The data presented here provide evidence that *A. phagocytophilum* infection leads to modified host cell gene transcription and phenotype by epigenetically altering host chromatin in regions that play a regulatory role in gene expression – a global mechanism for control of eukaryotic host cell function by intracellular bacteria. Other bacteria that closely interact with host cells, whether intracellular or not, may use similar mechanisms for manipulating host cell function. Further study of this host control mechanism could define prokaryotic effectors associated with this process and facilitate development of new strategies for the prevention and treatment of infections caused by bacteria with intimate host cell associations.

# **Materials and Methods**

#### Cell lines and cell culture

The acute monocytic leukemia THP-1 (ATCC CCL-240) cell line was used in this study because it supports *A. phagocytophilum* growth and can be readily transfected with siRNAs and plasmids as described below. THP-1 cells were grown in RPMI medium containing 10% FBS in a humidified incubator at 37°C with 5%  $CO_2$ . Cell density was kept  $<5\times10^5$  cells/mL by diluting with fresh medium every three days. The cell permeable HDAC inhibitors TSA and sodium butyrate were used at concentrations of 10–800 nM and 1–10 mM, respectively. The proportion of infected cells was determined by microscopic examination of LeukoStat-stained cells.

#### Anaplasma phagocytophilum culture and isolation

A GFP-expressing HGE-1strain of *A. phagocytophilum* [30], as well as the Webster, Slovenia and HZ strains were used in this study. Infected cells were grown until >90% of the cells were infected. Uninfected THP-1 cells were used to adjust the infection level to 10% as needed. To isolate cell-free *A. phagocytophilum*,  $2 \times 10^7$  infected cells were collected by centrifugation at  $500 \times \text{g}$  for 5 min. Bacteria were released by lysis using a 25%G syringe needle. Nuclei and cell debris were removed by low speed centrifugation (1,000×g, 5 min) and *A. phagocytophilum* organisms were collected from the supernatant by centrifugation at 14,000×g for 10 min. The bacterial preparation was finally resuspended in RPMI medium for *in vitro* infection of THP-1 cells at a multiplicity of infection (MOI) of 10 bacteria per cell. Infection levels were determined by microscopy examination of stained cells, by qPCR or by flow cytometry as described below.

#### Histone extraction and analysis

Histones were isolated from *A. phagocytophilum*-infected and uninfected cells by acid extraction as described previously [31]. Briefly, nuclei from  $2 \times 10^7$  cells were obtained by hypotonic lysis. Histones were then extracted with 0.4 N H<sub>2</sub>SO<sub>4</sub>, TCAprecipitated and washed with acetone. To study changes in histone posttranslational modifications, equal amounts (1 µg) of isolated histones were separated in 15% SDS-PAGE and analyzed by immunoblotting using antibodies specific for Ac-H3 and Me-H3.

#### Cell fractionation and immunoblotting

Nuclear extracts from  $2 \times 10^7$  *A. phagocytophilum*-infected or uninfected THP-1 cells were prepared using NE-PER Nuclear and Cytoplasmic Extraction Reagents (Pierce, USA) with protease inhibitors (PMSF and cOmplete, Roche, USA). Samples of the nuclear fractions were analyzed by immunoblotting to determine the presence of HDAC1 or HDAC2. Cytoplasmic fractions were used for normalization for  $\beta$ -actin content.

Approximately 1  $\mu$ g total protein was electrophoresed in a 10% or 15% SDS-PAGE gel and transferred to a nitrocellulose membrane. The blot was blocked with 3% bovine serum albumin, and probed with 1:2,000 HDAC1 rabbit polyclonal antibody (Sigma, USA), 4  $\mu$ g/mL HDAC2 monoclonal antibody (Sigma, USA), 1:100 Ac-H3 rabbit polyclonal antibody or 1:100 anti-Me-H3 (Millipore, USA). Human  $\beta$ -actin was used for normalization. Membranes were incubated with goat anti-mouse or anti-rabbit alkaline phosphatase conjugate (KPL, USA) and developed using Immun-Star AP substrate (Bio-Rad, USA). Band intensities were determined by densitometry using the public domain, free software ImageJ, Image processing and analysis in Java (http:// rsb.info.nih.gov/ij/).

## HDAC activity assay

Nuclear extracts from infected or uninfected THP-1 cells were prepared using NE-PER Nuclear and Cytoplasmic Extraction Reagents (Pierce, USA) as described above, and used to determine HDAC activity with the fluorescent HDAC assay kit (Active Motif,



**Figure 4. HDAC inhibition impairs the ability of** *A. phagocytophilum* **to propagate intracellularly.** THP-1 cells were infected with cell-free *A. phagocytophilum* and incubated with or without the HDAC inhibitors TSA and sodium butyrate. Twenty-four hours post infection, cells were collected and *A. phagocytophilum* infection level was determined by qPCR. The first point represents infection level in the absence of HDAC inhibitor. Cells were incubated with (A) TSA or (B) sodium butyrate. Infection levels were normalized for  $\beta$ -actin gene content to account for differences in the number of viable cells. (C) Cell-free *A. phagocytophilum* preincubated with 100 nM TSA for 2 hours prior to infection was used to infect THP-1 cells as described. *A. phagocytophilum* load was determined by qPCR 24 hours later. doi:10.1371/journal.ppat.1000488.g004



**Figure 5. HDAC1 silences defense gene expression and facilitates** *A. phagocytophilum* **infection of THP-1 cells.** (A) THP-1 cells were transfected with *HDAC1* or *HDAC2* siRNA and were infected 24 hours after transfection with cell-free *A. phagocytophilum*. The *A. phagocytophilum* load was determined 24 h after infection by qPCR and normalized to infection levels in non-transfected cells. (B) THP-1 cells were transfected with *HDAC1* siRNA for silencing or with pHDAC1-FLAG plasmid to overexpress HDAC1 and were infected 24 hours after transfection with cell-free GFP-expressing HGE-1 strain of *A. phagocytophilum*. The percent of infected cells was determined 24 h after infection by flow cytometry and normalized to infection levels in untreated cells. (C) Expression of defense genes by THP-1 cells infected with *A. phagocytophilum*–infected or transfected with pHDAC1-FLAG plasmid or *HDAC1* siRNA was determined by qRT-PCR. Gene expression was normalized to housekeeping genes and to the level of expression of untreated, uninfected THP-1 cells. doi:10.1371/journal.ppat.1000488.g005

USA) following the manufacturer's instructions. Nuclear extracts were diluted 1:3 in assay buffer and activity expressed as pmoles of fluorescent product formed after incubation at 37°C for 1 hour. Fluorescence intensity was measured using a Perkin Elmer Victor II plate reader with an excitation wavelength at 355 nm and emission at 460 nm.

#### RNA isolation and quantitative RT-PCR

To study the level of gene expression, total RNA from approximately  $2 \times 10^6$  cells was purified using RNeasy RNA extraction Kit (Qiagen, USA). cDNA was synthesized using SuperScript Reverse Transcriptase Kit (Invitrogen, USA) and quantitative PCR analysis was performed using primer sets specific for genes encoding for *HDAC1*, *HDAC2* or for defense genes using a custom-built PCR array (Table S1). Real time PCR was performed using SYBR Green Supermix in an iQ5 Multicycler (Bio-Rad, USA). Primer sequences were taken from the qPrimer-Depot database [32]. Expression levels were calculated using the comparative Ct method with the average Ct of human housekeeping genes *ACTB*, *B2M*, *GAPDH*, *HPRT1* and *RPL13A* as normalizer and uninfected cells as reference. The average±std. dev. of three experiments was calculated.

#### Chromatin immunoprecipitation

Analysis of histone modification patterns in the 1,000 bp proximal promoter regions of defense genes was performed using ChIP followed by qPCR using promoter-specific primers. Infected THP-1 cells  $(10^8)$  were cross-linked with formaldehyde at 1% final concentration for 15 min at room temperature. Cross-linking and chromatin isolation were performed as described [33]. Chromatin was sheared 4 times for 15 sec using a Branson Sonifier 250 (Branson, USA) at 1.5 constant output power. Histone-bound DNA was immunoprecipitated using Ac-H3 or Me-H3 -specific antibodies (Millipore, US). Input chromatin was normalized and uninfected THP-1 cells were used as control. A negative, isotypematched antibody control ChIP was also included. A sample of total chromatin was used as a positive control and for normalizing. Immunoprecipitated DNA fragments were quantitated by qPCR using primer sets specific for the defense gene promoter regions. The relative enrichment of each DNA fragment was calculated from the difference of the Ct with respect to the negative antibody control ChIP and normalized to the total chromatin control. Experiments were repeated two times and the average of the two determinations was calculated.

#### Plasmid and small interference RNA transfection

To overexpress HDAC1,  $2 \times 10^6$  THP-1 cells were nucleofected with 1 µg of pHDAC1-FLAG (Addgene plasmid 13820) purified using EndoFree Plasmid isolation kit (Qiagen, USA). Nucleofection was performed using Amaxa's nucleofector technology (Amaxa, Germany), Cell Line Nucleofector Kit V and program U-01. Cells transfected with plasmid vector pcDNA3.1 were used as control. Six hours after transfection, cells were infected with cell-free GFP-expressing *A. phagocytophilum* to assess the effect of HDAC1 overexpression on infection by flow-cytometry, expressed as changes in the proportion of infected THP-1 cells with respect to the control.

HDAC1 expression knockdown using siRNA, which has previously been demonstrated [6], was achieved using Stealth siRNA technology (Invitrogen, USA). RNAi conditions were optimized using Invitrogen's Green BLOCK-iT Fluorescent Oligo. THP-1 cells  $(5 \times 10^5 \text{ cells})$  were transfected with 200 pmole of each of 3 HDAC1 Stealth Select siRNA (HSS104725, HSS104726, HSS104727; Invitrogen, USA) or HDAC2 Stealth Select siRNA (HSS104728, HSS104729, HSS104730; Invitrogen, USA) using Lipofectamine RNAiMAX (Invitrogen, USA) in Opti-MEM. Negative Control siRNA #1 (Ambion, USA) was used as control. Twenty four hours after transfection, cells were infected with cell-free GFP-expressing *A. phagocytophilum* (MOI 10:1) and samples were taken at 24 hours post-infection for determination of HDAC silencing effect on *A. phagocytophilum* infection by qPCR or by flow-cytometry as described above.

#### Statistical methods

Statistical analysis was carried out using 2-sided Student's t-test for comparison means, where a p value <0.05 was considered significant. Error bars used throughout indicate standard error of the mean.

#### **Supporting Information**

Figure S1 HDAC expression and activity is increased during infection with different isolates of A. phagocytophilum. HL-60 cells were infected with Webster, Slovenia and HZ strains of A. phagocytophilum for 48 hours. (A) Changes in the amount of HDAC1 with infection were determined by immunoblotting using an HDAC1-specific antibody. β-actin was used as a control. HDAC1 band intensity was determined by densitometric analysis and HDAC1 protein expression level changes with respect to uninfected cells were calculated. Samples were normalized for βactin content. (B) HL-60 cells (HL) uninfected or infected with Webster (W), Slovenia (S) and HZ strains (HZ) of A. phagocytophilum were incubated for 24 h with and without 400 nM TSA. Histones were acid-extracted and acetylated and methylated H3 were detected by immunoblotting. Changes in the H3 acetylation and methylation patterns were determined by densitometric analysis of the immunoblot bands, and expressed as fold-change after TSA treatment with respect to the corresponding untreated control. Found at: doi:10.1371/journal.ppat.1000488.s001 (0.44 MB TIF)

Table S1 Genes and primers used in this study.

Found at: doi:10.1371/journal.ppat.1000488.s002 (0.03 MB DOC)

#### Acknowledgments

The authors would like to thank Dr. Dennis J. Grab for fruitful discussions and critical reading of the manuscript and Emily Clemens for technical assistance. We thank Dr. Eric M. Verdin for plasmid pHDAC1-FLAG, Miroslav Petrovec for the Slovenia-1 strain of *A. phagocytophilum*, Yasuko Rikihisa for the HZ strain, and Roderick Felsheim, Michael Herron and Ulrike Munderloh for the GFP-expressing HGE-1 strain. We also thank Dr. John Audia for helpful comments and critical evaluation of the results.

#### **Author Contributions**

Conceived and designed the experiments: JCGG JSD. Performed the experiments: JCGG NCB SJT. Analyzed the data: JCGG SJT JSD. Contributed reagents/materials/analysis tools: JCGG NCB SJT JSD. Wrote the paper: JCGG JSD. Designed and did most of the experiments and prepared the manuscript: JCGG. Helped with cell culture model of infection, qPCRs and analysis of human isolates: NCB. Was involved in the HDAC inhibitor studies: SJT.

- Galan JE, Cossart P (2005) Host-pathogen interactions: a diversity of themes, a variety of molecular machines. Current Opinion in Microbiology 8: 1–3.
- Bryant PA, Venter D, Robins-Browne R, Curtis N (2004) Chips with everything: DNA microarrays in infectious diseases. Lancet Infectious Diseases 4: 100–111.
- Zupkovitz G, Tischler J, Posch M, Sadzak I, Ramsauer K, et al. (2006) Negative and positive regulation of gene expression by mouse histone deacetylase 1. Mol Cell Biol 26: 7913–7928.
- 4. Glaser KB, Staver MJ, Waring JF, Stender J, Ulrich RG, et al. (2003) Gene expression profiling of multiple histone deacetylase (HDAC) inhibitors: defining a common gene set produced by HDAC inhibition in T24 and MDA carcinoma cell lines. Mol Cancer Ther 2: 151–163.
- Lee HS, Park MH, Yang SJ, Jung HY, Byun SS, et al. (2004) Gene expression analysis in human gastric cancer cell line treated with trichostatin A and Sadenosyl-L-homocysteine using cDNA microarray. Biol Pharm Bull 27: 1497–1503.
- Glaser KB, Li J, Staver MJ, Wei RQ, Albert DH, et al. (2003) Role of class I and class II histone deacetylases in carcinoma cells using siRNA. Biochem Biophys Res Commun 310: 529–536.
- Valls E, Blanco-Garcia N, Aquizu N, Piedra D, Estaras C, et al. (2007) Involvement of chromatin and histone deacetylation in SV40 T antigen transcription regulation. Nucleic Acids Res 35: 1958–1968.
- Punga T, Akusjarvi G (2000) The adenovirus-2 E1B-55K protein interacts with a mSin3A/histone deacetylase 1 complex. FEBS Lett 476: 248–252.
- Radkov SA, Touitou R, Brehm A, Rowe M, West M, et al. (1999) Epstein-Barr virus nuclear antigen 3C interacts with histone deacetylase to repress transcription. J Virol 73: 5688–5697.
- Hamon MA, Cossart P (2008) Histone modifications and chromatin remodeling during bacterial infections. Cell Host Microbe 4: 100–109.
- Hamon MA, Batsche E, Regnault B, Tham TN, Seveau S, et al. (2007) Histone modifications induced by a family of bacterial toxins. Proc Natl Acad Sci U S A 104: 13467–13472.
- Arbibe L, Kim DW, Batsche E, Pedron T, Mateescu B, et al. (2007) An injected bacterial effector targets chromatin access for transcription factor NF-kappaB to alter transcription of host genes involved in immune responses. Nat Immunol 8: 47–56.
- Mott J, Rikihisa Y, Tsunawaki S (2002) Effects of Anaplasma phagocytophila on NADPH oxidase components in human neutrophils and HL-60 cells. Infection and Immunity 70: 1359–1366.
- Scaife H, Woldehiwet Z, Hart CA, Edwards SW (2003) Anaplasma phagocytophilum reduces neutrophil apoptosis in vivo. Infection and Immunity 71: 1995–2001.
- Garyu JW, Choi KŠ, Grab DJ, Dumler JS (2005) Defective phagocytosis in Anaplasma phagocytophilum-infected neutrophils. Infect Immun 73: 1187–1190.
- Choi KS, Grab DJ, Dumler JS (2004) Anaplasma phagocytophilum infection induces protracted neutrophil degranulation. Infect Immun 72: 3680–3683.
- Kim HY, Rikihisa Y (2000) Expression of interleukin-1beta, tumor necrosis factor alpha, and interleukin-6 in human peripheral blood leukocytes exposed to human granulocytic ehrlichiosis agent or recombinant major surface protein P44. Infect Immun 68: 3394–3402.

- Akkoyunlu M, Fikrig E (2000) Gamma interferon dominates the murine cytokine response to the agent of human granulocytic ehrlichiosis and helps to control the degree of early rickettsemia. Infect Immun 68: 1827–1833.
- de la Fuente J, Ayoubi P, Blouin EF, Almazan C, Naranjo V, et al. (2005) Gene expression profiling of human promyelocytic cells in response to infection with *Anaplasma phagocytophilum*. Cellular Microbiology 7: 549–559.
- Borjesson DL, Kobayashi SD, Whitney AR, Voyich JM, Argue CM, et al. (2005) Insights into pathogen immune evasion mechanisms: *Anaplasma phagocytophilum* fails to induce an apoptosis differentiation program in human neutrophils. J Immunol 174: 6364–6372.
- Pedra JH, Sukumaran B, Carlyon JA, Berliner N, Fikrig E (2005) Modulation of NB4 promyclocytic leukemic cell machinery by *Anaplasma phagocytophilum*. Genomics 86: 365–377.
- Lee HC, Kioi M, Han J, Puri RK, Goodman JL (2008) Anaplasma phagocytophiluminduced gene expression in both human neutrophils and HL-60 cells. Genomics 92: 144–151.
- Garcia-Garcia JC, Rennoll-Bankert KE, Pelly S, Milstone AM, Dumler JS (2009) Silencing of Host Cell CYBB Gene Expression by the Nuclear Effector AnkA of the Intracellular Pathogen Anaplasma phagocytophilum. Infect Immun.
- Park J, Kim KJ, Choi KS, Grab DJ, Dumler JS (2004) Anaplasma phagocytophilum AnkA binds to granulocyte DNA and nuclear proteins. Cell Microbiol 6: 743–751.
- Caturegli P, Asanovich KM, Walls JJ, Bakken JS, Madigan JE, et al. (2000) ankA: an Ehrlichia phagocytophila group gene encoding a cytoplasmic protein antigen with ankyrin repeats. Infect Immun 68: 5277–5283.
- Carlyon JÅ, Chan WT, Galan J, Roos D, Fikrig E (2002) Repression of rac2 mRNA expression by *Anaplasma phagocytophila* is essential to the inhibition of superoxide production and bacterial proliferation. J Immunol 169: 7009–7018.
- Wong ET, Jenne DE, Zimmer M, Porter SD, Gilks CB (1999) Changes in chromatin organization at the neutrophil elastase locus associated with myeloid cell differentiation. Blood 94: 3730–3736.
- Lu Y, Goldenberg I, Bei L, Andrejic J, Eklund EA (2003) HoxA10 represses gene transcription in undifferentiated myeloid cells by interaction with histone deacetylase 2. J Biol Chem 278: 47792–47802.
- Wang Y, Curry HM, Zwilling BS, Lafuse WP (2005) Mycobacteria inhibition of IFN-gamma induced HLA-DR gene expression by up-regulating histone deacetylation at the promoter region in human THP-1 monocytic cells. J Immunol 174: 5687–5694.
- Felsheim RF, Herron MJ, Nelson CM, Burkhardt NY, Barbet AF, et al. (2006) Transformation of *Anaplasma phagocytophilum*. BMC Biotechnol 6: 42.
- Shechter D, Dormann HL, Allis CD, Hake SB (2007) Extraction, purification and analysis of histones. Nat Protoc 2: 1445–1457.
- Cui W, Taub DD, Gardner K (2007) qPrimerDepot: a primer database for quantitative real time PCR. Nucleic Acids Res 35: D805–809.
- Ren B, Robert F, Wyrick JJ, Aparicio O, Jennings EG, et al. (2000) Genomewide location and function of DNA binding proteins. Science 290: 2306–2309.